

**Delta Coal**  
**Manning & CVC Collieries**

**Lake Macquarie Benthos Survey**

**Results No. 18**



**By Dr Emma Laxton**

**August 2020**

**J.H. & E.S. Laxton - Environmental Consultants P/L**  
**170 Warrimoo Avenue, St. Ives. NSW 2075 Australia**  
**Telephone: 0429 855 891 Email: [emmalaxton07@gmail.com](mailto:emmalaxton07@gmail.com)**

## Table of Contents

1.	Summary of survey findings	5
2.	Introduction	8
3.	Location of Sampling Stations	9
4.	Selection and Evaluation of the Sampling Method	9
5.	Sampling Procedure	15
6.	Factors Affecting the Depth of water in Lake Macquarie	16
7.	Water Quality of Lake Macquarie (April 1983 – March 1997)	18
8.	Light attenuation in Lake Macquarie (1983 – 1997)	19
9.	Results	20
10.	Benthos of Study Area – February 2012 to August 2020	21
11.	Molluscs found as dead shells	27
12.	Benthic organisms of the Study Area – August 2020	34
13.	Sediment Analysis	46
14.	Water Quality Profiles – August 2020	48
15.	Analysis of data	50
16.	Conclusions	53
17.	References and Acknowledgements	54

## List of Tables

1.	Co-ordinates of Benthos Sampling Stations prepared by the LDO team	13
2.	Water Quality of the body of Lake Macquarie (1983-1997)	18
3.	Organisms found at sampling Stations on 15 <sup>th</sup> and 18 <sup>th</sup> August 2020	35
4.	Number of species found at each station from February 2012 to August 2020	43

5.	Mean number of marine benthic organisms for Control, Reference and Impact Stations (August 2020)	45
6.	Percent mud in sediment from each station - February 2012 to August 2020	46
7.	Description of sediments from each Sampling Station collected in August 2020	47
8.	Physical characteristics of the bottom water – August 25 <sup>th</sup> 2020	49

## **List of Figures**

1a	Location of Benthos Sampling Stations (August 2020)	11
1b	Extent of mining to March 2020	12
2.	Water level changes in a coastal lagoon with an entrance open to coastal waters	17
3.	Mean percentage changes in PAR with depth at station 1 – Wyee Point over 12 months	19
4.	Actual percentage change in PAR at Station 1 – Wyee Point on the morning of 21-4-83	20
5.	Relationship between marine benthic organisms and stations	52

## **List of Plates**

1a.	Annelid species found in benthos of Lake Macquarie (February 2012 – August 2020)	22
1b.	Gastropod species found in benthos of Lake Macquarie (February 2012 – August 2020)	23
1c.	Bivalve species found in benthos of Lake Macquarie (February 2012 – August 2020)	24
1d.	Brittle stars are found amongst the mussel beds of Lake Macquarie, NSW	27
1e.	Crab species found in Lake Macquarie, NSW	27
2a.	Large shell removed from samples during sorting process – February 2012 survey	28
2b.	Small shell removed from samples during sorting process – February 2012 survey	28
2c.	Large shells removed from samples – September 2012 survey	29
2d.	Small shells removed from samples – September 2012 survey	29

2e.	Large shells removed from samples during sorting in March 2013	30
2f.	Small shells removed from samples during sorting in March 2013	30
3a.	Mollusc species found as dead shells in the benthos of Lake Macquarie, NSW	32
3b.	Gastropod species found as dead shells in the benthos of Lake Macquarie, NSW	34

## **Appendix 1**

Water Quality Profiles for the August 2020 marine benthos survey, Lake Macquarie	55
--	----

## Summary of survey findings

In August 2020, 22 benthic stations were sampled. The following is a history of benthos sampling from 2014 to 2020:

- By March 2014, mining beneath Lake Macquarie had proceeded so that two Reference stations (R) had been re-designated Impact Stations (IM), namely:

R3 became IM5

R4 became IM6.

- By September 2014, Station R5 had become the impact station IM7.
- In March 2016 two more stations were added to the sampling schedule. They were:

C5 GR 367701 6334310

R7 GR 366232 6333856.

- In September 2016, difficult geology beneath Bardens Bay and along parts of Summerland Point led Lake Coal to begin mining beneath Chain Valley Bay. To accommodate this change in mining direction, three additional benthos sampling stations were added. They were C6, R8 and R9.

C6 GR 363988 6332492

R8 GR 364523 6332010

R9 GR 365258 6331210

- The total number of Stations sampled in September 2017 was 19.
- In March 2018, three new stations were added to the sampling programme. They were:

C7 GR 366276 6334947

R10 GR 365172 6334706

R11 GR 367072 6333639

- The mud basin off Summerland Point, in Chain Valley Bay and Bardens Bay, was found to be inhabited by 24 species of organisms greater than 1mm in size. This list was derived from the 18 samplings undertaken between February 2012 and August 2020. Polychaete worms and bivalve molluscs were the most frequently encountered animals.
- Bottom sediment in the study area was composed of fine black mud with varying proportions of

black sand and shell fragments.

Water levels in Lake Macquarie can vary by as much as 1.3m over the course of a year due to combinations of the following phenomena:

- Diurnal tidal changes (around 0.05m)
- Changes in atmospheric pressure (up to 0.4m)
- Wave set up at the entrance to Lake Macquarie
- Inflow of water from the catchment during major rainfall events.

Light attenuation through the water column of Lake Macquarie, measured off Wyee Point, between 1983 and 1985, showed that only 14% of the photosynthetically active radiation (PAR) reached the lake bed at 2m depth (the growth limit of seagrasses and macroscopic algae in the LDO study area). At 6m depth, between 2% and 4% of the surface PAR reached the lakebed, not enough light to support the growth of seagrasses or benthic algae.

The 18 samplings of the benthos undertaken at six monthly intervals between February 2012 and August 2020 revealed the following:

- The same suite of organisms dominated each of the 22 sample stations. These were polychaete worms and bivalves.
- Stations were distinguished by the relative abundance of the dominant species.
- Water depth was not in any way important in determining the species composition at a station.
- Physical variables such as salinity, conductivity and turbidity of the bottom water had little influence on the species composition of the benthos. Dissolved oxygen concentration, however, can have a major effect on abundance. It is clear that major extinction events have occurred in the mud basin of Lake Macquarie. The evidence for this lies in the presence of large numbers of intact but dead bivalve shells entombed in the mud. The cause of extinction events appears to be prolonged dissolved oxygen depletion of bottom water. Prolonged dissolved oxygen depletion of the bottom water was measured during the water quality study conducted by Laxton and Laxton (1983 to 1997). Low concentrations of dissolved oxygen in the bottom water were also recorded during the March 2020 sampling period. Stations with low abundance of organisms correlated with low concentrations in the bottom waters.
- In March and September 2019, the total number of organisms found in sediment from the 22 stations was 832 and 815, respectively. This was around half the number of organisms collected in September 2018 (1576 organisms). In March and August 2020, a total of 1032 and 1367

organisms were collected. It appears an extinction event has occurred relatively recently in Lake Macquarie, although the increased numbers of organisms collected in the 2020 studies suggest the lake is in recovery.

These results appear to support the notion that increasing the water depth by the predicted 0.8m subsidence has, to date, had no discernible effect on the composition and abundance of organisms making up the benthos of the mud basin.

Up until recently, little significant rain has fallen in the catchments of Lake Macquarie. Annual rainfall in the Cooranbong (Lake Macquarie AWS) region was 839.8 mm in 2017; 859.8 mm in 2018 and 763.4 mm in 2019 (BOM Station Number 061412). The lack of rainfall caused the salinity of the water column to become very high (over 39 parts per thousand by March 2019) and almost uniform from surface to bottom. The Lake Macquarie region has since received heavy rainfall in August (111.2 mm) and September (64.8 mm) 2019 and again in January (79.6 mm), February (335.4 mm), March (173.0 mm) and July (184.0 mm) 2020 (BOM Station Number 061412). This rainfall lowered the salinity of water in the lake to around 36 parts per thousand in 2019, 33 parts per thousand in March 2020, and 32 parts per thousand in August 2020. Water temperature, salinity, conductivity and pH were found to be uniform from surface to bottom. During the August 2020 sampling period, dissolved oxygen concentrations were above 100% saturation throughout the water profile and study area.

The lack of rainfall also influenced water clarity. The water of the lake became very clear for long periods. This high water clarity led to some interesting effects on the benthos of the study area. First, the small seagrass, *Halophila sp.* became established as a dense bed in 6m of water at Station R10 (Brightwaters Bay) in September 2018. *Halophila sp.* was not recorded at Station R10 in March 2019 but in September 2019 a healthy plant of *Zostera capricorni* was found at this station. Second, at stations C4 and IM2, red and brown algae were found on mussels at depths between 4.5 and 6 m of water in September 2018.

In September 2019 some changes to the composition of the upper 100mm of the bottom sediments were detected. At Stations C1-C4 and C6-C7 no sand was present, just fine black silt. This indicated that these sediments had been reworked since March 2019. Sediments at Stations R5, R6 R8 and R9 also appeared to have been reworked. In March 2020, changes were again detected. Sediments at stations C5 and C7 comprised mostly of coarse black sand. In August 2020, sediment comprised mostly of fine black silt.

Note: AWS - Automatic Weather Station

## Introduction

In 2012 Lake Coal P/L was seeking a variation to its mining agreement because of proposed changes to its mining methods. They were planning on increasing minwall panel widths to 85m wide, 97m total extraction, which will result in some additional subsidence above that currently approved. As such, a modification and supporting EA was prepared. The predicted subsidence agreed to by the NSW Government was around 0.406m. The method now proposed will increase subsidence to around 0.468m.

NSW Department of Planning and Infrastructure raised concerns that this increase in depth of water over the existing benthic community of the mud basin of Lake Macquarie may alter the species composition and relative abundances of organisms within that community.

To address these concerns, Lake Coal decided to conduct a benthic survey of the mud basin community to attempt to answer the following questions.

- What is the structure of the benthic community of the mud basin off Summerland Point and in Chain Valley Bay?
- What changes to the benthic community, if any, have taken place in areas of the lake mud basin that have been subjected to subsidence from previous mining activity?
- What changes to the benthic community, if any, may be expected in the mud basin community from the proposed variation to the mining method?

This study had a seasonal component and the benthos could change from year to year without the influence of any subsidence due to mining.

Ms Jemma Sargent of JSA Environmental prepared a formal document entitled:

Benthic Communities Management Plan. Chain Valley Colliery Domains 1 & 2 Continuation. Project (10\_0161). 25 June 2012.

The extraction plan required under Condition 6 of Schedule 3 within Project approval (10\_0161) requires that a Benthic Communities Management Plan (BCMP) be developed. This BCMP was prepared to provide for the management of the potential impacts and/or environmental consequences of the proposed second workings on benthic communities and includes:

- surveys of the lakebed to enable contours to be produced and changes in depth following subsidence to be accurately measured
- benthic species surveys within the area subject to second workings, as well as control sites outside the area subject to second workings (at similar depths) to establish baseline data on species numbers and composition within the communities

- a program of ongoing seasonal monitoring of benthic species in both control and impact sites
- development of a model to predict the likely impact of increased depth and associated subsidence impacts and effects, including but not limited to light reduction and sediment disturbance, on benthic species number and benthic communities composition, incorporating the survey data collected.

Three types of station were sampled. They were:

- Control stations – C, areas of lakebed sufficiently remote from previous or proposed mining.
- Reference stations – R, areas of lakebed above subsidence areas of previous mining.
- Impact stations – IM areas of lakebed where subsidence is expected from future mining.

Two depth zones within the mud basin were sampled, -4.5m AHD and -5.5 to -6.0m AHD. The locations of the sampling stations were specified by Mr Chris Ellis and Mr Wade Covey, using the results of a bathymetric survey of the lake and the known locations of past and proposed mining.

In November 2014, project consent 10\_0161 was surrendered. It was replaced by consent SSD-5465 as modified. The remodeled subsidence values were 0.62m for the single seam extraction area (everywhere that is currently being mined) and up to 0.886m for areas where multiseam mining will occur (near Site R2).

This report (August 2020) presents the results of the just completed 18<sup>th</sup> sampling of the now 22 (previously 19, 16, 14 and 12) stations off Summerland Point, in Chain Valley Bay, Bardens Bay and Sugar Bay. These results will be compared with those obtained from the previous seventeen surveys (February 2012 to March 2020). The benthic survey was conducted between the 15<sup>th</sup> and 18<sup>th</sup> August 2020. Water quality variables were measured on 25<sup>th</sup> August 2020. The work in August 2020 was supervised by Mr Lachlan McWha of Chain Valley Colliery.

## Location of Sampling Stations

**Figure 1** shows the location of sampling stations, depth contours of the lake, and the locations of existing and proposed underground mine workings prepared by Mr Chris Armit and the LDO team in February 2017 and updated in October 2019. **Table 1** provides the exact location of each sampling station by latitude and longitude and by eastings and northings using WGS84 datum.

## Selection and Evaluation of the Sampling Method

Methods for sampling benthos of sedimentary bottoms of oceans, coastal waters and saline and

freshwater lakes must fulfill the following criteria:

- The area of bottom collected by the sampling device must be appropriate to the types and sizes of organisms inhabiting the substratum.
- The depth that the sampler penetrates the sediment must be sufficient to capture infauna or identifiable parts of more deeply buried species.
- Sufficient samples must be taken within the benthic environment to be certain that more than 95% of the component species of the ecosystem are collected.
- Sufficient samples must be taken to permit the population densities of component species to be calculated.

In 1971, Dr John Laxton was appointed by Dr Frank Talbot, Director of the Australian Museum, to lead a team of scientists to undertake the Shelf Benthic Survey. The purpose of the Shelf Benthic Survey was to provide baseline biological data on the benthos, fish and birds of the Continental Shelf adjacent to Sydney. Baseline oceanographic and biological data for coastal waters adjacent to Sydney were required to evaluate the effects of the proposed deep water ocean outfalls planned by the Metropolitan Water Sewerage and Drainage Board to replace the existing shoreline sewage outfalls. Both rocky bottoms and sedimentary bottoms were present in the study area and water depths ranged from the intertidal zone out to 200m.




Figure 1a. Location of Benthos Sampling Stations (August 2020).



Figure 1b. Extent of mining to March 2020

**Table 1.** Co-ordinates of Benthos Sampling Stations prepared by the LDO team.

Station Co-ordinate Table					
Station	Sample depth (m) AHD	Latitude	Longitude	MG-56 Easting	MG56 Northing
C1	-4.50	S33° 09' 10.69"	E151° 32' 50.11"	364519	6330815
C2	-4.50	S33° 08' 02.89"	E151° 33' 56.65"	366214	6332927
C3	-5.50	S33° 07' 55.78"	E151° 33' 49.05"	366014	6333144
C4	-6.00	S33° 08' 06.35"	E151° 32' 41.17"	364260	6332794
C5	-6.00			367701	6334310
C6	-5.50			363988	6332492
C7	-5.50			366276	6334947
IM1	-4.50	S33° 09' 13.44"	E151° 32' 58.51"	364738	6330734
IM2	-4.50	S33° 08' 24.67"	E151° 33' 03.34"	364842	6332237
IM3	-5.50	S33° 08' 29.02"	E151° 32' 57.52"	364693	6332101
IM4	-6.00	S33° 08' 09.42"	E151° 32' 57.04"	364873	6332705
R1	-4.50	S33° 08' 47.18"	E151° 32' 37.31"	364177	6331535
R2	-4.50	S33° 09' 28.23"	E151° 33' 43.87"	365919	6330294
R3 (IM5)	-5.50	S33° 08' 00.10"	E151° 32' 56.72"	364660	6332992
R4 (IM6)	-6.00	S33° 08' 07.58"	E151° 33' 00.88"	364771	6332763
R5(IM7)	-5.50	S33° 07' 30.78"	E151° 32' 40.55"	364229	6333889
R6	-6.00	S33° 07' 22.56"	E151° 32' 52.42"	364533	6334146
R7	-6.00			366232	6333856
R8	-5.50			364523	6332010

R9	-4.50			365258	6331210
R10	-5.50			365172	6334706
R11	-6.00			367072	6333639

At first, a Shipek grab was employed to collect samples of sediment. The Shipek grab used a spring loaded hemi-cylindrical bucket that rotated through 180° to collect a half cylinder of sediment nominally 200 x 200 mm in area and cut to a maximum depth of 100mm. On gravel bottoms, the Shipek grab worked consistently to collect 200 x 200 x 100mm samples. On sandy bottoms the grab, when triggered, penetrated the bottom to varying depths, collecting half cylinders of sediment that could range in depth from the full 100mm to as little as 25mm. This meant that the area of the seabed sampled varied greatly between samples taken at the same station and the depth of some samples was so shallow that many species of infauna were not collected. On muddy bottoms the heavy Shipek grab could plunge into the soft mud and emerge untriggered.

The Shipek grab was safe to use from a pitching and rolling vessel but as a scientific sampling device, it had serious deficiencies.

The Shelf Benthic Survey then obtained a Smith-MacIntyre (S-M) grab for evaluation. The Smith-MacIntyre grab used two spring operated clam-shells which swung inwards towards the midline to gather 200 x 200 x 100mm samples of sediment. This grab also had similar limitations to the Shipek grab when used to sample various sediment types. The worst feature of the S-M grab was that the two springs had to be tensioned by a lever separately and then a keeper was placed in position to stop it triggering while on deck or while being lowered to the sea bed. To position the keeper, the operator had to reach in between the two cocked spring loaded clam-shells. These clam-shell jaws were sharp and the action was violent enough to remove a hand. The Captain of the vessel banned its use on the project and undoubtedly saved someone's hand.

Following completion of the Shelf Benthic Survey, John Laxton joined an engineering firm that was commissioned to design wastewater outfalls for Gosford City, Wyong Shire and the Hunter Area. Baseline data on water quality and biology were again required and the seabed in the discharge and mixing zones was either rocky or sedimentary. As the maximum water depth in sedimentary areas was 30m, diver operated sample collection devices could be used to sample sedimentary bottoms. It was decided to build a diver operated benthic sampler that would overcome the difficulties and deficiencies of the available grab samplers. It should collect a 200 x 200 x 100mm section of sediment consistently and be easy to operate in conditions of zero underwater visibility.

To collect a 200 x 200 x 100mm sample of sediment consistently an aluminium box was designed that

could be slid sideways into the sediment, whether gravel, sand or mud, and be filled completely before it was lifted clear of the bottom and the door closed and locked to retain the sediment. The top of the box included a panel of 1.0mm stainless steel mesh. Thus each box contained its own sieve to permit particles less than 1mm in size to be removed from the box leaving only large particles and organisms.

Tests of this box revealed that in all sediments (gravel, sand and mud) between 3 and 5 replicate samples were required to capture 95% of the species present. Once the maximum number of replicates required had been determined, five sieve boxes were manufactured along with a carry case to contain the boxes on the journey between the surface and the bottom and back. These devices permitted samples of consistent area and depth to be collected. Five replicates were always collected regardless of the sediment type or the environment being studied so that individual species/area curves were not required for each new area being investigated.

Five sieve boxes sample an area of 0.20 m<sup>2</sup>. This sampling device has been used in all J.H. & E.S. Laxton - Environmental Consultants P/L benthos studies since 1980.

In an attempt to make the Summerland Point/Chain Valley Bay study results comparable with other studies, the BCMP required two cores of 100mm diameter and 200mm depth to be taken along with the 5 sieve box samples. These two cores covered an area of 0.015 m<sup>2</sup>. There was no requirement in the BCMP to determine how many cores of these dimensions were needed to capture 95% of the benthic species inhabiting the lakebed. However, it is unlikely that sampling 0.015 m<sup>2</sup> of bottom sediment will provide a more realistic picture of the structure of the benthic community than sampling 0.20 m<sup>2</sup> of bottom sediment.

## **Sampling Procedure**

Between September 2012 and August 2020, five replicate samples of basin mud were collected at each station using 200 x 200 x 100mm sieve boxes (1mm mesh). Two 100 x 200mm core samples were also collected at each station on each date sampled.

Twenty-two stations were sampled in August 2020. At each station the following procedure was carried out:

- A GPS unit was used to locate the sampling station. The boat was positioned upwind of the station and was then allowed to drift back to the exact location. When the wind strength was 0-5km/h, the boat stayed on position. When the wind strength increased from 5 to 25km/h, the boat yawed on its anchor warp, causing the distance from the boat to the station to vary greatly and the sampling difficulty to increase. This was mitigated by working in calm conditions only.
- A line with five sieve boxes, two 100 x 200mm core samplers and a mesh bag containing a 250mL jar for whole sediment was cast over board as the boat drifted into position.

- The diver descended to the lakebed to fill the 250mL jar, the two core samplers and five sieve boxes with sediment.
- The samplers were then hauled to the surface, and the contents of each sampler placed in a clean, labeled zip-lock plastic bag.
- Processing of samples occurred in the laboratory.
- A water quality profile from surface to bottom (at 0.5m depth intervals) was taken using a calibrated Yeo-Kal 618 Water Quality Analyser. Water temperature, conductivity, salinity, pH, dissolved oxygen, turbidity and depth were measured. Each line of data was stored in the memory of the machine.

In the laboratory the marine benthic samples were treated in the following way:


- Each sample was tipped into a 1 mm mesh sieve and washed free of mud.
- The washed material from each sample was then placed into an enamel dish and sorted for animals.
- Organisms and parts of organisms were removed, counted, identified and the results entered into a spread sheet. This process was repeated until the debris of the entire sample had been examined.
- Sorted organisms were preserved in formaldehyde solution.
- All shell remaining in the sample was kept for later examination.

The 250mL samples of whole sediment were treated in the following way:

- Each sample was tipped into a 1L clear glass measuring cylinder and the volume made up to 600mL with seawater.
- The cylinders were stoppered and shaken vigorously to suspend the sediment in the seawater.
- The cylinders were then placed on the laboratory bench to allow the fractions of the sediment to settle.
- Once settled the sediment profiles were photographed and the volumes of each fraction (shell and coarse sand, fine sand, mud and fine silt) were calculated and recorded. Results were displayed relative to the initial volume of sediment collected in the 250mL jar.

## **Factors Affecting the Depth of Water in Lake Macquarie**

The bathymetric chart of Lake Macquarie shows water depths relative to AHD. The actual depth of water above the lakebed varied greatly (**Figure 2**).



**Figure 2.** Water level changes in a coastal lagoon with an entrance open to coastal waters.

The actual water depth above the lakebed varied between 0 and 1.3m above AHD over a year. Water depths in coastal saline lakes with an open entrance to coastal waters varied due to combinations of the following factors:

- The body of Lake Macquarie was subject to tidal influence. The height of the tidal prism at Swansea Head may reach almost 2m (during spring tides) but by the time the body of the lake was reached, the tidal prism had been reduced to around 0.05m.
- The height of coastal waters and coastal lakes were influenced by changes in atmospheric pressure. The Tasman Sea acted as a huge barometer. When the atmospheric pressure was high the sea surface was depressed. This caused water to drain from Lake Macquarie causing the depth of water in the body of the lake to decrease. When the atmospheric pressure over the Tasman Sea was low, the surface of the sea bulged upwards. This raising of sea level caused water to flow into Lake Macquarie, increasing the water depth.
- Low pressure systems in the Tasman Sea almost always generated strong winds and coastal rainfall. The strong winds caused large swells to form that impact the coast. Wave setup at the entrance to Lake Macquarie caused the water level in the lake to rise as large volumes of seawater entered the system.
- Rainfall during a period of low atmospheric pressure caused runoff into catchment rivers and streams to increase. When this extra water reached the body of Lake Macquarie, the water level rose in proportion to the runoff volume. This water was prevented from exiting the lake by wave setup at the entrance and the state of the tide. Under these circumstances, the level of the lake

rose to heights of a meter or more above AHD (**Figure 2**).

## Water Quality of Lake Macquarie (April 1983 – March 1997)

In 1983 the Hunter District Water Board (later Hunter Water Corporation) commissioned J.H. & E.S. Laxton – Environmental Consultants P/L to carry out a water quality study of Lake Macquarie in conjunction with their plans to sewer the western shore of the lake. The study commenced in April 1983 with monthly sampling of the lake and ended in March 1997. The water quality results for the body of Lake Macquarie (as opposed to the creeks) are summarized and presented in **Table 2**.

**Table 2.** Water Quality of the body of Lake Macquarie (1983-1997)

Variable		Mean	Maximum	Minimum
<b>Water Temperature (°C)</b>	Surface	20.56	33.77	10.95
	Bottom	20.06	29.17	11.45
<b>Water Salinity (ppt)</b>	Surface	32.61	37.96	1.00
	Bottom	33.92	37.95	21.06
<b>PH</b>	Surface	8.28	9.28	7.19
	Bottom	8.26	8.90	7.55
<b>Dissolved Oxygen (% saturation)</b>	Surface	101.6	177.7	71.9
	Bottom	89.5	147.0	0.9
<b>Turbidity (NTU)</b>	Surface	3.0	32.8	0.0
	Bottom	5.1	77.7	0.0
<b>Transmission of light through water (%)</b>	Surface	94.2	99.9	7.3
	Bottom	88.1	99.4	2.0
<b>Total Suspended Solids (mg/L)</b>	Surface	4.8	123.5	0.5
<b>Chlorophyll-a (µg/L)</b>	Surface	2.953	112.900	0.000
<b>Ammonia-nitrogen (mg-N/L)</b>	Surface	0.071	1.500	0.006
	Bottom	0.075	0.813	0.010
<b>Organic-nitrogen (mg-N/L)</b>	Surface	0.355	9.691	0.000
	Bottom	0.361	3.357	0.002
<b>Oxidized-nitrogen (mg-N/L)</b>	Surface	0.010	0.459	0.000
	Bottom	0.008	0.142	0.000


<b>Total-nitrogen (mg-N/L)</b>	Surface	0.436	9.749	0.033
	Bottom	0.445	3.918	0.027
<b>Orthophosphate phosphorus (mg-P/L)</b>	Surface	0.0191	0.4148	0.0006
	Bottom	0.0188	0.1386	0.0003
<b>Total phosphorus (mg-P/L)</b>	Surface	0.0450	0.8922	0.0025
	Bottom	0.0489	0.3534	0.0022
<b>Faecal coliform bacteria (No./100mL)</b>	Surface	55	5000	0

Blue shading in Table 2 indicates variables of interest to this study of the benthos of Lake Macquarie.


## Light attenuation in Lake Macquarie (1983 – 1997)

Observations made over many years (Laxton, 2007) show that photosynthetic benthic organisms (seagrasses and algae) are confined to the shallow water areas around the perimeter of Lake Macquarie. In Chain Valley Bay, Bardens Bay and off Summerland Point, seagrasses and benthic algae grow between 0 and -1.89m below AHD (except in September 2018 when *Halophila* and some algae were found in 4.5 to 6m of water at some stations due to low rainfall and clear water).

The water quality study of Lake Macquarie, carried out between 1983 and 1997, measured Photosynthetically Active Radiation (PAR) changes with depth monthly at twelve stations throughout the lake during the years 1983 to 1985. Data for Station 1 off Wyee Point are presented in **Figure 3** and **Figure 4**.



**Figure 3.** Mean percentage changes in PAR with depth at Station 1 - Wyee Point over 12 months.



**Figure 4.** Actual percentage changes in PAR at Station 1 - Wyee Point on the morning of 21-4- 83.

It was found that only 14% of the light present at the surface reached a depth of 2.0m below the surface. By 6m below the surface only between 2% and 4% of PAR remained. Seagrasses and algae just manage to survive at 14% of the surface radiation but have no chance of survival at 6m below the surface. The mud basin of Lake Macquarie was devoid of macroscopic benthic algae and seagrasses except at some stations in September 2018.

## Results

### Benthos of the Study Area – February 2012 to August 2020

The following organisms were found in the sediment samples collected off Summerland Point and in Chain Valley Bay between February 2012 and August 2020:

Designated name	Family or Species	Comments
Anemone	Coelenterata	Found associated with mussel shells.
Planaria (Flat worm)	Platyhelminthes	2 specimens found in 2017.
Polychaete thin	<i>Sthenelais pettiboneae</i>	Most common polychaete present.
Polychaete (thick)	Cirratulidae	Present in small numbers.
Polychaete (mud tube)	Not yet identified	Present in small numbers.
Polychaete	Terebellidae	Present at Stations C1, C6, R1 and IM2.
<i>Pectinaria sp.</i> Polychaete	Terebellidae	First found in March 2019
Gastropod	<i>Nassarius jonasii</i>	Present in small numbers.
Gastropod	<i>Lepsiella (Bedeva) hanleyi</i>	Present in small numbers.
Gastropod	Bullimorph slug	One specimen found in August 2014.

Bivalve	<i>Corbula truncata</i>	Common as live animals and dead shells.
Bivalve	<i>Soletellina alba</i>	Common
Bivalve	<i>Paphia undulata</i>	Uncommon as live animals. Common as dead shells.
Bivalve	<i>Cyamiomactra mactroides</i>	Uncommon. (Brown or pink bivalve)
Bivalve	<i>Anadara trapezia</i>	Uncommon.
Bivalve	<i>Dosinia sculpta</i>	Many juveniles found in sandy sediment in September 2019.
Bivalve	<i>Trichomya hirsuta</i>	Common as dead shells. Found in large clumps at C2, C6, R3, R7, IM2 and IM3.
Ophuroid	Brittle star	Uncommon, found associated with mussel clumps and on mud.
Sponge	White calcareous sponge	Specimen found associated with mussels.
	Pink sponge	Small species found on mud surface.
	Red sponge	Several specimens found in 2019.
Crab	Small	Uncommon.
Prawn	Small	One specimen taken in March 2013 at R3 and one specimen in September 2013 at C4.
Shrimp	Small	Found at IM2 in March 2014.
Fish	Small (35mm)	One specimen taken at C3 (September 2012), at R1 (September 2013) and at IM4 in March 2017. 1 specimen in C6 in 2019.

**Plates 1a to 1d** provide information about the benthic organisms present in the basin mud of Lake Macquarie, NSW.

**Plate 1a.** Annelid species found in the benthos of Lake Macquarie (February 2012 - August 2020).



**Phylum:** Annelida  
**Class:** Polychaeta  
**Subclass:** Errantia  
**Order:** Phyllodocida  
**Family:** Sigalionidae  
**Genus:** *Sthenelais*  
**Species:** *Sthenelais pettiboneae*

**Remarks:** Found in marine environments.



**Phylum:** Annelida  
**Class:** Polychaeta  
**Subclass:** Canalipalpata  
**Order:** Terebellida  
**Family:** Cirratulidae

**Remarks:** Cirratulids vary in size from 1-20 cm long. They are mostly burrowers in soft sediments but some live in rock crevices. The head is conical or wedge-shaped and has no antennae. The body is generally cylindrical, tapering at both ends. Cirratulids are characterised by a large number of simple elongate filaments along the body. The genera are poorly defined.

**Plate 1b.** Gastropod species found in the benthos of Lake Macquarie (February 2012 - August 2020).



**Phylum:** Mollusca  
**Class:** Gastropoda  
**Superfamily:** Buccinoidea  
**Family:** Nassariidae  
**Genus:** *Nassarius*  
**Species:** *Nassarius jonasii*

**Remarks:** Endemic to Australia; Noosa Heads, Qld, to SA. Inhabit sand and mud flats in estuaries and lagoons, intertidal down to 100 m. Most *Nassarius* species are very active scavengers. They often burrow into marine substrates and then wait with only their siphon protruding, until they smell nearby food.



**Phylum:** Mollusca  
**Class:** Gastropoda  
**Order:** Neogastropoda  
**Family:** Muricidae  
**Genus:** *Lepsiella* (*Bedevea*)  
**Species:** *Lepsiella hanleyi*

**Remarks:** Common name mussel drill. Shell up to 32 mm, with angulated whorls, a high spire and moderately long anterior canal and with both spiral threads and axial ribs. Endemic to Australia. Found in temperate and southern parts of tropical Australia. Lives mainly on sheltered shores, including estuaries and often in association with mangroves. Feeds by drilling holes in bivalves. Lays lens-shaped capsules and development is direct.

**Plate 1c.** Bivalve species found in the benthos of Lake Macquarie (February 2012 - August 2020).



**Phylum:** Mollusca  
**Class:** Bivalvia  
**Order:** Myoida  
**Family:** Corbulidae  
**Genus:** *Corbula*  
**Species:** *Corbula truncata*

**Remarks:** Marine bivalve mollusc.



**Phylum:** Mollusca  
**Class:** Bivalvia  
**Order:** Veneroida  
**Family:** Psammobiidae  
**Genus:** *Soletellina*  
**Species:** *Soletellina alba*

**Remarks:** Posterior and anterior margins almost parallel. Shell thin and normally bluish, rarely white. Lives intertidally and subtidally in sand and mud, especially in sheltered environments. Occurs all around Australia; not recorded elsewhere.



**Phylum:** Mollusca  
**Class:** Bivalvia  
**Order:** Veneroida  
**Family:** Veneridae  
**Genus:** *Paphia*  
**Species:** *Paphia undulata*

**Remarks:** Saltwater clam, marine bivalve mollusc. Inhabits inshore shallow sandy seabeds.



**Phylum:** Mollusca  
**Class:** Bivalvia  
**Order:** Veneroida  
**Family:** Veneridae  
**Genus:** *Dosinia*  
**Species:** *Dosinia sculpta*

**Remarks:** *Dosinia* is a genus of saltwater clams, marine bivalve molluscs in the family Veneridae, (subfamily Dosiniinae). The shell of *Dosinia* species is disc-like in shape, usually white, and therefore is reminiscent of the shells of Lucinid bivalves.

Typically found in the intertidal zone at the water's edge at a mean distance from sea level of -15 meters (-50 feet).



**Phylum:** Mollusca  
**Class:** Bivalvia  
**Order:** Veneroida  
**Family:** Cyamiidae  
**Genus:** *Cyamiomactra*  
**Species:** *Cyamiomactra mactroides*



**Phylum:** Mollusca  
**Class:** Bivalvia  
**Order:** Arcoida  
**Family:** Arcidae  
**Genus:** *Anadara*  
**Species:** *Anadara trapezia*

**Remarks:** Sydney cockle, or ark cockle is an estuarine filter-feeding bivalve. Its calcareous, heavily-ribbed, shell can grow to approximately 7 to 8 cm across. Its current range is along the east coast of Australia, from Queensland to Victoria. It has been used as an indicator species to study levels of the metals selenium, copper and cadmium.



**Phylum:** Mollusca  
**Class:** Bivalvia  
**Order:** Mytiloida  
**Family:** Mytilidae  
**Genus:** *Trichomya*  
**Species:** *Trichomya hirsuta*

**Remarks:** The hairy mussel is a major part of the megafauna of Lake Macquarie. It is tolerant of low oxygen levels in the water and its temperature tolerance range has been researched in connection with using the waters of the lake for cooling power stations.

Hairy mussels have been used as bioindicators to monitor concentrations of heavy metals (namely Pb, Cd, Cu, Zn, Co, Ni, and Ag) in marine environments.

**Plate 1d.** Brittle stars are found amongst the mussel beds of Lake Macquarie, NSW.



**Phylum:** Echinodermata  
**Class:** Ophiuroidea  
**Order:** Ophiurida  
**Family:** Ophionereididae  
**Genus:** *Ophionereis*  
**Species:** *Ophionereis schayeri*

**Remarks:** Largest and most common brittle star found in Sydney waters. Brittle stars have five long, slender arms which radiate out from a central disc. The mouth is located in the centre of the underside of the disc. There is no anus. Offshore, brittle stars form dense aggregations. In intertidal zones, they are typically found as single individuals in crevices, under stones and amongst seaweed. They feed by raising their arms above the substrate; extending tube-feet; and removing particles from the water. They pass food along the arms to the mouth. They also scavenge on decaying matter. They inhabit the hairy mussel beds of Lake Macquarie.

**Plate 1e** Crab species found in Lake Macquarie, NSW



**Phylum:** Arthropoda  
**Class:** Malacostraca  
**Order:** Decapoda

## Molluscs found as dead shells

Benthic organism samples collected between February 2012 and August 2020 included a large

component of shell. **Plates 2a** and **2b** show the mass of shell obtained from the sixty 200x200x100mm samples of sediment taken in February 2012. **Plate 2c** and **Plate 2d** show the mass of shell collected in September 2012 and **Plates 2e** and **2f** show the mass of shells collected in March 2013.



**Plate 2a.** Large shell removed from samples during sorting process - February 2012 survey.



**Plate 2b.** Small shells removed from samples during sorting process - February 2012 survey.



**Plate 2c.** Large shells removed from samples - September 2012 survey.



**Plate 2d.** Small shells removed from samples during sorting in September 2012.



**Plate 2e.** Large shells removed from samples during sorting in March 2013.



**Plate 2f.** Small shells removed from samples during sorting in March 2013.

Similar masses of shell were found in the samples of the September 2013 to August 2020 surveys. These masses of shell were photographed for the record but were not included in this report.

The following molluscs were found in the large volume of shell collected during the sampling periods between February 2012 and August 2020:

- |                                |  |
|--------------------------------|--|
| 1. <i>Paphia undulata</i>      | 7. <i>Chlamys</i> sp.                          |
| 2. <i>Anomia</i> sp.           | 8. <i>Saccostrea glomerata</i>                 |
| 3. <i>Dosinia sculpta</i>      | 9. <i>Corbula truncata</i>                     |
| 4. <i>Trichomya hirsuta</i>    | 10. <i>Batillaria (Velacumantis) australis</i> |
| 5. <i>Katelysia rhytiphora</i> | 11. <i>Conuber</i> sp.                         |
| 6. <i>Pecten</i> sp.           | 12. <i>Anadara trapezia</i>                    |

**Plates 3a** and **3b** provide information about bivalve mollusc and gastropod species found as dead shells in the basin mud of Lake Macquarie, New South Wales during the periods of sampling.

**Plate 3a.** Mollusc species found as dead shells in the benthos of Lake Macquarie, NSW.



**Phylum :** Mollusca

**Class:** Bivalvia

**Order:** Ostreoida

**Family:** Anomiidae

**Genus:** *Anomia*

**Remarks:** Genus of saltwater clam, marine bivalve mollusc. Known as "jingle shells". Common in both tropical and temperate oceans and live primarily attached to rock or other shells via a calcified byssus that extends through the lower valve. *Anomia* shells tend to take on the surface shape of what they are attached to; thus if an *Anomia* is attached to a scallop shell, the shell of the *Anomia* will also show ribbing.



**Phylum :** Mollusca

**Class:** Bivalvia

**Order:** Veneroida

**Family:** Veneridae

**Genus:** *Katelysia*

**Species:** *Katelysia rhytiphora*

**Remarks:** Commonly known as mud cockles, this group of commercially important bivalves often represents a major faunal component of shallow estuarine and marine embayments. *K. rhytiphora* is broadly distributed around Australia's temperate coastline from Augusta, Western Australia to Port Jackson, NSW.



**Phylum :** Mollusca

**Class:** Bivalvia

**Order:** Ostreoida

**Family:** Pectinidae

**Genus:** *Pecten*

**Remarks:** Genus of large saltwater clams or scallops. Marine bivalve mollusc.



**Phylum :** Mollusca

**Class:** Bivalvia

**Order:** Ostreoida

**Family:** Pectinidae

**Genus:** *Chlamys*

**Remarks:** Genus of saltwater clams or scallops. Marine bivalve mollusc.



**Phylum :** Mollusca

**Class:** Bivalvia

**Order:** Ostreoida

**Family:** Pectinidae

**Genus:** *Saccostrea*

**Species:** *Saccostrea glomerata*

**Remarks:** Sydney rock oysters are endemic to Australia and New Zealand. In Australia it is found in bays, inlets and sheltered estuaries from Wingan Inlet in eastern Victoria, along the east coast of NSW and up to Hervey Bay QLD, around northern Australia and down the west coast to Shark Bay in WA. Sydney rock oysters are capable of tolerating a wide range of salinities (halotolerant). They are usually found in the intertidal zone to 3 metres (9.8 ft) below the low water mark.

**Plate 3b.** Gastropod species found as dead shells in the benthos of Lake Macquarie, NSW.



**Phylum :** Mollusca  
**Class:** Gastropoda  
**Family:** Naticidae  
**Genus:** *Conuber*  
**Species:** *Conuber sordidum*

**Remarks:** Species of predatory sea snail. A marine gastropod mollusc known commonly as the moon snail. Lives on intertidal muddy sand flats near mangroves or sea weed.



**Phylum :** Mollusca  
**Class:** Gastropoda  
**Family:** Batillariidae  
**Genus:** *Batillaria (Velacumantis)*  
**Species:** *Batillaria australis*

**Remarks:** The Australian Mud Whelk is a marine gastropod found on mud flats in estuaries, river mouths and mangrove swamps. The snail has a high resistance to predation and environmental tolerance, which may partially explain its success as an invasive species. This species is one of the hosts for the flatworm parasite *Austrobilharzia*. Larvae of the flatworm are discharged from the snail into the surrounding water. They normally burrow into the legs of wading birds and complete their life cycle, but may burrow through the skin of humans, causing "bathers itch".

## **Benthic organisms in the Study Area - August 2020**

The organisms found living in the sediments of the mud basin off Summerland Point and in Chain Valley Bay and Bardens Bay were entered into an Excel worksheet. **Table 3** shows the organisms found in each replicate at each station sampled in August 2020. Data for sieve box samples were separated from data obtained from core samples.

**Table 3. Organisms found at Sampling Stations on 15<sup>th</sup> and 18<sup>th</sup> August 2020.**

<b>Control Station C1</b>		Depth -4.50m AHD			56 364519		6330815		Sampled August 15th-18th 2020							
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab	
C1.1	2	1	0	0	0	2	16	0	0	1	0	0	0	0	0	
C1.2	1	0	0	0	0	6	9	0	0	0	0	0	0	0	0	
C1.3	2	0	0	0	0	3	4	0	0	0	0	1	0	0	0	
C1.4	1	3	0	0	0	2	15	0	0	0	0	0	0	0	0	
C1.5	2	0	0	0	0	4	22	0	0	0	0	0	0	0	0	
C1.6	0	0	0	0	0	1	5	0	0	0	0	0	0	0	0	
C1.7	0	0	0	0	0	2	7	0	0	0	0	0	0	0	0	
Mean/station (boxes)	1.6	0.8	0.0	0.0	0.0	3.4	13.2	0.0	0.0	0.2	0.0	0.2	0.0	0.0	0.0	
Mean cores	0.0	0.0	0.0	0.0	0.0	1.5	6.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
no./m2 (box)	40	20	0	0	0	85	330	0	0	5	0	5	0	0	0	
no./m2 (core)	0	0	0	0	0	30	120	0	0	0	0	0	0	0	0	
<b>No. species (box)</b>	<b>6</b>															
<b>No. species (core)</b>	<b>2</b>															
<b>Total Organisms at Station</b>															<b>97</b>	
<b>Control Station C2</b>		Depth -4.50m AHD			56 366214		6332927		Sampled August 15th-18th 2020							
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Planaria	
C2.1	5	3	0	0	0	9	5	0	0	0	0	0	0	0	0	
C2.2	0	8	0	0	0	2	13	0	0	1	0	0	0	0	0	
C2.3	3	6	0	0	0	2	5	0	0	0	0	0	0	0	0	
C2.4	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
C2.5	6	3	0	0	0	4	10	0	0	0	0	0	0	0	0	
C2.6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
C2.7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mean/station (boxes)	3.0	4.0	0.0	0.0	0.0	3.4	6.6	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	
Mean cores	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
no./m2 (box)	75	100	0	0	0	85	165	0	0	5	0	0	0	0	0	
no./m2 (core)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
<b>No. species (box)</b>	<b>5</b>															
<b>No. species (core)</b>	<b>0</b>															
<b>Total Organisms at Station</b>															<b>86</b>	
<b>Control Station C3</b>		Depth -5.50m AHD			56 366014		6333144		Sampled August 15th-18th 2020							
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Bivalve Dosinia	Planaria	Sponge	
C3.1	2	0	0	0	0	0	1	0	0	0	0	0	0	0	0	
C3.2	4	1	0	0	0	6	8	0	0	0	0	0	0	0	0	
C3.3	3	0	0	0	0	0	4	0	0	0	0	0	0	0	0	
C3.4	3	2	0	0	0	3	4	0	0	0	0	0	0	0	0	
C3.5	2	0	0	0	0	3	0	0	0	0	0	0	0	0	0	
C3.6	1	0	0	0	0	0	3	1	0	0	0	0	0	0	0	
C3.7	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mean/station (boxes)	2.8	0.6	0.0	0.0	0.0	2.4	3.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Mean cores	0.5	1.0	0.0	0.0	0.0	0.0	1.5	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
no./m2 (box)	70	15	0	0	0	60	85	0	0	0	0	0	0	0	0	
no./m2 (core)	10	20	0	0	0	0	30	10	0	0	0	0	0	0	0	
<b>No. species (box)</b>	<b>4</b>															
<b>No. species (core)</b>	<b>4</b>															
<b>Total Organisms at Station</b>															<b>46</b>	

<b>Control Station C4</b>		Depth -5.50m AHD			56 364260		6332794		Sampled August 15th-18th 2020								
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab		
C4.1	1	0	0	0	0	5	25	0	0	0	0	0	0	0	0		
C4.2	2	2	0	0	0	0	24	0	0	0	0	0	0	0	0		
C4.3	3	1	0	0	0	0	5	0	0	0	0	0	0	0	0		
C4.4	1	0	0	0	0	3	17	0	0	0	0	0	0	0	0		
C4.5	1	1	0	0	0	6	14	0	0	0	0	0	0	0	0		
C4.6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
C4.7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Mean/station (boxes)	1.6	0.8	0.0	0.0	0.0	2.8	17.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
Mean cores	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
no./m2 (box)	40	20	0	0	0	70	425	0	0	0	0	0	0	0	0		
no./m2 (core)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
<b>No. species (box)</b>	<b>4</b>																
<b>No. species (core)</b>	<b>0</b>																
<b>Total Organisms at Station</b>															<b>111</b>		
<b>Control Station C5</b>		Depth -5.50m AHD			56 367701		6334510		Sampled August 15th-18th 2020								
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab		
C5.1	5	5	0	0	0	0	0	0	0	0	0	3	0	0	0		
C5.2	3	13	0	0	0	0	0	0	0	0	0	0	0	0	0		
C5.3	8	3	0	0	0	0	0	0	0	0	0	1	0	0	0		
C5.4	5	13	0	0	0	0	0	0	0	0	0	4	0	0	0		
C5.5	0	5	0	0	0	0	0	0	0	0	0	0	0	0	0		
C5.6	1	5	0	0	0	0	0	0	0	0	0	0	0	0	0		
C5.7	2	2	0	0	0	0	0	0	0	0	0	1	0	0	0		
Mean/station (boxes)	4.2	7.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.6	0.0	0.0	0.0		
Mean cores	1.5	3.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0		
no./m2 (box)	105	195	0	0	0	0	0	0	0	0	0	40	0	0	0		
no./m2 (core)	30	70	0	0	0	0	0	0	0	0	0	10	0	0	0		
<b>No. species (box)</b>	<b>3</b>																
<b>No. species (core)</b>	<b>3</b>																
<b>Total Organisms at Station</b>															<b>68</b>		
<b>Control Station C6</b>		Depth -5.50m AHD			56 363988		6332492		Sampled August 15th-18th 2020								
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab		
C6.1	2	0	0	0	0	1	0	0	0	0	0	0	0	0	0		
C6.2	0	0	0	0	0	1	5	0	0	0	0	0	0	0	0		
C6.3	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
C6.4	2	0	0	0	0	0	5	0	0	1	0	0	0	0	0		
C6.5	2	1	0	0	0	4	4	0	0	0	0	0	0	0	0		
C6.6	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
C6.7	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0		
Mean/station (boxes)	1.4	0.2	0.0	0.0	0.0	1.2	2.8	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0		
Mean cores	0.5	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
no./m2 (box)	35	5	0	0	0	30	70	0	0	5	0	0	0	0	0		
no./m2 (core)	10	0	0	0	0	0	10	0	0	0	0	0	0	0	0		
<b>No. species (box)</b>	<b>5</b>																
<b>No. species (core)</b>	<b>2</b>																
<b>Total Organisms at Station</b>															<b>29</b>		

Control Station C7		Depth -5.50m AHD			56 364736		6334947		Sampled August 15th-18th 2020											
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab					
C7.1	4	2	0	0	0	0	0	0	0	0	0	0	1	0	0	0				
C7.2	0	1	0	0	0	0	2	0	0	0	0	0	1	0	0	0				
C7.3	4	2	0	0	0	1	0	0	0	0	0	0	0	0	0	0				
C7.4	2	6	0	0	0	0	0	0	0	0	0	0	1	0	0	0				
C7.5	2	5	0	0	0	0	0	0	0	0	0	0	1	0	0	0				
C7.6	1	1	0	0	0	0	1	0	0	0	0	0	1	0	0	0				
C7.7	1	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
Mean/station (boxes)	2.4	3.2	0.0	0.0	0.0	0.2	0.4	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.0	0.0				
Mean cores	1.0	2.5	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.8	0.0	0.0	0.0				
no./m2 (box)	60	80	0	0	0	5	10	0	0	0	0	0	20	0	0	0				
no./m2 (core)	20	50	0	0	0	0	10	0	0	0	0	0	16	0	0	0				
No. species (box)	5																			
No. species (core)	4																			
<b>Total Organisms at Station</b>															<b>35</b>					
Station R1		Depth -4.50m AHD			56 364177		6331535		Sampled August 15th-18th 2020											
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab					
R1.1	2	0	0	0	0	1	6	0	0	0	0	0	0	0	0	0				
R1.2	1	2	0	0	0	1	11	0	0	0	0	0	0	0	0	0				
R1.3	2	0	0	0	0	1	16	0	0	0	0	0	0	0	0	0				
R1.4	3	2	0	0	0	2	3	0	0	0	0	0	0	0	0	0				
R1.5	1	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0				
R1.6	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0				
R1.7	0	0	0	0	0	0	2	0	0	0	0	0	1	0	0	0				
Mean/station (boxes)	1.8	0.8	0.0	0.0	0.0	1.0	7.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0				
Mean cores	0.5	0.0	0.0	0.0	0.0	0.5	1.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0				
no./m2 (box)	45	20	0	0	0	25	190	0	0	0	0	0	0	0	0	0				
Mean cores	10	0	0	0	0	10	20	0	0	0	0	0	10	0	0	0				
No. species (box)	4																			
No. species (core)	4																			
<b>Total Organisms at Station</b>															<b>56</b>					
Station R2		Depth -4.50m AHD			56 365919		6330294		Sampled August 15th-18th 2020											
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab					
R2.1	1	1	0	0	0	6	8	0	0	0	0	0	0	0	0	0				
R2.2	3	1	0	0	0	0	11	0	0	1	0	0	0	0	0	0				
R2.3	0	1	0	0	0	9	6	0	0	1	0	0	0	0	0	0				
R2.4	1	0	0	0	0	6	6	0	0	0	0	0	0	0	0	0				
R2.5	3	0	0	0	0	4	3	0	0	0	0	0	0	0	0	0				
R2.6	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0				
R2.7	2	0	0	0	0	1	5	0	0	0	0	0	0	0	0	0				
Mean/station (boxes)	1.6	0.6	0.0	0.0	0.0	5.0	6.8	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0				
Mean cores	1.0	0.0	0.0	0.0	0.0	1.0	3.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0				
no./m2 (box)	40	15	0	0	0	125	170	0	0	10	0	0	0	0	0	0				
no./m2 (core)	20	0	0	0	0	20	60	0	0	0	0	0	0	0	0	0				
No. species (box)	5																			
No. species (core)	3																			
<b>Total Organisms at Station</b>															<b>72</b>					

Station R3 (now IM5)		Depth -5.50m AHD			56 364660		6332992		Sampled August 15th-18th 2020								
Replicates		Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Bivalve Saccostrea	Ophuroid	Fish	Crab	
R3.1	IM5.1	2	1	0	0	0	0	2	0	0	0	15	0	0	0	0	
R3.2	IM5.2	2	0	0	0	0	0	0	0	0	0	24	1	0	0	0	
R3.3	IM5.3	1	1	0	0	0	0	8	1	0	0	0	0	0	0	0	
R3.4	IM5.4	4	0	0	0	0	0	3	0	0	0	28	0	0	0	0	
R3.5	IM5.5	2	0	0	0	0	1	2	0	0	0	22	0	0	0	0	
R3.6	IM5.6	0	0	0	0	0	2	2	0	0	0	1	0	0	0	0	
R3.7	IM5.7	2	2	0	0	0	1	2	0	0	0	0	0	0	0	0	
Mean/station (boxes)		2.2	0.4	0.0	0.0	0.0	0.2	3.0	0.2	0.0	0.0	17.8	0.2	0.0	0.0	0.0	
Mean cores		1.0	1.0	0.0	0.0	0.0	1.5	2.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	
no./m2 (box)		55	10	0	0	0	5	75	5	0	0	445	5	0	0	0	
no./m2 (core)		20	20	0	0	0	30	40	0	0	0	10	0	0	0	0	
<b>No. species (box)</b>		<b>7</b>															
<b>No. species (core)</b>		<b>5</b>															
<b>Total Organisms at Station</b>																<b>132</b>	
Station R4 (now IM6)		Depth -6.00m AHD			56 364771		6332763		Sampled August 15th-18th 2020								
Replicates		Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab	
R4.1	IM6.1	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	
R4.2	IM6.2	1	1	0	0	0	0	1	0	0	0	0	0	0	0	0	
R4.3	IM6.3	1	1	0	0	0	0	2	0	0	0	0	0	0	0	0	
R4.4	IM6.4	2	2	0	0	0	4	10	0	0	0	0	0	0	0	0	
R4.5	IM6.5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
R4.6	IM6.6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
R4.7	IM6.7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mean/station (boxes)		1.0	0.8	0.0	0.0	0.0	1.0	2.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Mean cores		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
no./m2 (box)		25	20	0	0	0	25	65	0	0	0	0	0	0	0	0	
no./m2 (core)		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
<b>No. species (box)</b>		<b>4</b>															
<b>No. species (core)</b>		<b>0</b>															
<b>Total Organisms at Station</b>																<b>27</b>	
Station R5 (now IM7)		Depth -6.00m AHD			56 364229		6333889		Sampled August 15th-18th 2020								
Replicates		Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab	
R5.1		3	0	0	0	0	5	3	0	0	1	0	0	0	0	0	
R5.2		1	0	2	0	0	4	1	1	0	0	0	0	0	0	0	
R5.3		1	1	0	0	0	3	0	0	0	1	0	0	0	0	0	
R5.4		2	0	0	0	0	1	2	0	0	0	0	0	0	0	0	
R5.5		3	0	0	0	0	0	14	0	0	0	0	0	0	0	0	
R5.6		0	0	0	0	0	2	3	0	0	1	0	0	0	0	0	
R5.7		1	0	0	0	0	0	1	0	0	1	0	0	0	0	0	
Mean/station (boxes)		2.0	0.2	0.4	0.0	0.0	2.6	4.0	0.2	0.0	0.4	0.0	0.0	0.0	0.0	0.0	
Mean cores		0.5	0.0	0.0	0.0	0.0	1.0	2.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.0	
no./m2 (box)		50	5	10	0	0	65	100	5	0	10	0	0	0	0	0	
no./m2 (core)		10	0	0	0	0	20	40	0	0	20	0	0	0	0	0	
<b>No. species (box)</b>		<b>7</b>															
<b>No. species (core)</b>		<b>4</b>															
<b>Total Organisms at Station</b>																<b>49</b>	

Station R6 (now IM8)		Depth -6.00m AHD			56 364533		6334146		Sampled August 15th-18th 2020							
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab	
R6.1	2	1	0	0	0	5	17	0	0	0	0	0	0	0	0	
R6.2	0	0	0	0	0	4	22	0	0	0	0	0	0	0	0	
R6.3	4	0	1	0	0	2	6	0	0	0	0	0	0	0	0	
R6.4	3	1	0	0	0	2	10	0	0	0	0	0	0	0	0	
R6.5	1	0	0	0	0	3	11	0	0	0	0	0	0	0	0	
R6.6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
R6.7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mean/station (boxes)	2.0	0.4	0.2	0.0	0.0	3.2	13.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Mean cores	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
no./m2 (box)	50	10	5	0	0	80	330	0	0	0	0	0	0	0	0	
no./m2 (core)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
No. species (box)	5															
No. species (core)	0															
<b>Total Organisms at Station</b>														<b>95</b>		
Station R7		Depth -6.00m AHD			56 366232		6333856		Sampled August 15th-18th 2020							
Replicates	Polychaete thin	Polychaete mud	Polychaete Terebellid	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab	
R7.1	0	0	0	0	0	0	0	0	0	0	7	0	0	0	1	
R7.2	0	0	0	0	0	1	0	0	0	0	30	0	0	0	0	
R7.3	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0	
R7.4	3	2	0	0	0	0	9	0	0	0	0	0	0	0	0	
R7.5	4	3	0	0	0	0	5	0	0	1	0	0	0	0	0	
R7.6	1	0	0	0	0	1	1	0	0	0	0	0	0	0	0	
R7.7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mean/station (boxes)	1.4	1.0	0.0	0.0	0.0	0.2	2.8	0.0	0.0	0.2	7.4	0.6	0.0	0.0	0.2	
Mean cores	0.5	0.0	0.0	0.0	0.0	0.5	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
no./m2 (box)	35	25	0	0	0	5	70	0	0	5	185	15	0	0	5	
no./m2 (core)	10	0	0	0	0	10	10	0	0	0	0	0	0	0	0	
No. species (box)	8															
No. species (core)	3															
<b>Total Organisms at Station</b>														<b>69</b>		
Station R8		Depth -6.00m AHD			56 364323		63322010		Sampled August 15th-18th 2020							
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab	
R8.1	0	0	0	0	0	0	4	0	0	0	0	0	0	0	0	
R8.2	4	0	0	0	0	0	1	1	0	0	0	0	0	0	0	
R8.3	1	0	0	0	0	2	1	0	0	0	0	0	0	0	0	
R8.4	2	0	0	0	0	0	3	0	0	0	0	0	0	0	0	
R8.5	1	0	0	0	0	0	13	0	0	0	0	0	0	0	0	
R8.6	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	
R8.7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Mean/station (boxes)	1.6	0.0	0.0	0.0	0.0	0.4	4.4	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Mean cores	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
no./m2 (box)	40	0	0	0	0	10	110	5	0	0	0	0	0	0	0	
no./m2 (core)	0	0	0	0	0	0	10	0	0	0	0	0	0	0	0	
No. species (box)	4															
No. species (core)	1															
<b>Total Organisms at Station</b>														<b>33</b>		

<b>Station R9</b>		Depth -6.00m AHD			56 366232		6331210		Sampled August 15th-18th 2020								
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab		
R9.1	2	0	0	0	0	2	12	0	0	0	0	0	0	0	0		
R9.2	4	0	0	0	0	3	8	0	0	0	0	0	0	0	0		
R9.3	3	0	0	0	0	3	9	1	0	0	0	0	0	0	0		
R9.4	1	0	0	0	0	3	2	0	0	0	0	0	0	0	0		
R9.5	2	0	0	0	0	7	12	0	0	1	0	0	0	0	0		
R9.6	0	0	0	0	0	1	6	0	0	0	0	0	0	0	0		
R9.7	1	0	0	0	0	0	3	0	0	0	0	0	0	0	0		
Mean/station (boxes)	2.4	0.0	0.0	0.0	0.0	3.6	8.6	0.2	0.0	0.2	0.0	0.0	0.0	0.0	0.0		
Mean cores	0.5	0.0	0.0	0.0	0.0	0.5	4.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
no./m2 (box)	60	0	0	0	0	90	215	5	0	5	0	0	0	0	0		
no./m2 (core)	10	0	0	0	0	10	90	0	0	0	0	0	0	0	0		
<b>No. species (box)</b>	<b>5</b>																
<b>No. species (core)</b>	<b>3</b>																
													<b>Total Organisms at Station</b>		<b>75</b>		
<b>Station R10</b>		Depth -6.00m AHD			56 365172		6334708		Sampled August 15th-18th 2020								
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab		
R10.1	0	2	0	0	0	0	4	0	0	0	0	0	0	0	0		
R10.2	2	2	0	0	0	2	3	0	0	0	0	0	0	0	0		
R10.3	3	2	0	0	0	1	8	0	0	0	0	0	0	0	0		
R10.4	2	0	0	0	0	1	0	0	0	0	0	0	0	0	0		
R10.5	3	2	0	0	0	1	4	1	0	0	0	0	0	0	0		
R10.6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
R10.7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Mean/station (boxes)	2.0	1.6	0.0	0.0	0.0	1.0	3.8	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
Mean cores	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0		
no./m2 (box)	50	40	0	0	0	25	95	5	0	0	0	0	0	0	0		
no./m2 (core)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
<b>No. species (box)</b>	<b>5</b>																
<b>No. species (core)</b>	<b>0</b>																
													<b>Total Organisms at Station</b>		<b>43</b>		
<b>Station R11</b>		Depth -6.00m AHD			56 367072		6333638		Sampled August 15th-18th 2020								
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab		
R11.1	0	3	0	0	0	0	0	0	0	0	0	2	0	0	0		
R11.2	1	2	0	0	0	0	1	0	0	0	0	3	0	0	0		
R11.3	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0		
R11.4	4	6	0	0	0	0	0	0	0	0	0	1	0	0	0		
R11.5	1	4	0	0	0	0	2	0	0	0	0	1	0	0	0		
R11.6	2	1	0	0	0	0	0	0	0	0	0	1	0	0	0		
R11.7	3	2	0	0	0	1	0	0	0	0	0	2	0	0	0		
Mean/station (boxes)	1.4	3.2	0.0	0.0	0.0	0.0	0.6	0.0	0.0	0.0	0.0	1.4	0.0	0.0	0.0		
Mean cores	2.5	1.5	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	1.5	0.0	0.0	0.0		
no./m2 (box)	35	80	0	0	0	0	15	0	0	0	0	35	0	0	0		
no./m2 (core)	50	30	0	0	0	10	0	0	0	0	0	30	0	0	0		
<b>No. species (box)</b>	<b>4</b>																
<b>No. species (core)</b>	<b>4</b>																
													<b>Total Organisms at Station</b>		<b>33</b>		

<b>Station IM1</b>		Depth -4.50m AHD			56 364738		6330734		Sampled August 15th-18th 2020										
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Sponge	Dosinia				
IM1.1	3	1	0	0	0	4	2	0	0	0	0	0	0	0	0				
IM1.2	4	0	0	0	0	2	3	0	0	0	0	0	0	0	0				
IM1.3	3	0	0	0	0	2	6	0	0	1	0	0	0	0	0				
IM1.4	5	0	0	0	0	2	2	0	0	0	0	0	0	0	0				
IM1.5	0	0	0	0	0	3	3	0	0	0	0	0	0	0	0				
IM1.6	2	0	0	0	0	0	3	0	0	0	0	0	0	0	0				
IM1.7	1	2	0	0	0	0	4	0	0	0	0	0	0	0	0				
Mean/station (boxes)	3.0	0.2	0.0	0.0	0.0	2.6	3.2	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0				
Mean cores	1.5	1.0	0.0	0.0	0.0	0.0	3.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0				
no./m2 (box)	75	5	0	0	0	65	80	0	0	5	0	0	0	0	0				
no./m2 (core)	30	20	0	0	0	0	70	0	0	0	0	0	0	0	0				
<b>No. species (box)</b>	<b>5</b>																		
<b>No. species (core)</b>	<b>3</b>																		
<b>Total Organisms at Station</b>															<b>46</b>				
<b>Station IM2</b>		Depth -4.50m AHD			56 364842		6332237		Sampled August 15th-18th 2020										
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Sponge	Crab				
IM2.1	4	1	0	0	0	0	2	0	0	0	8	1	0	0	0				
IM2.2	1	0	0	0	0	0	0	0	0	0	30	0	0	0	1				
IM2.3	2	3	0	0	0	0	2	0	0	0	0	0	0	0	0				
IM2.4	1	3	0	0	0	0	6	0	0	0	0	0	0	0	0				
IM2.5	2	3	0	0	0	0	0	0	0	0	15	0	0	0	0				
IM2.6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
IM2.7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
Mean/station (boxes)	2.0	2.0	0.0	0.0	0.0	0.0	2.0	0.0	0.0	0.0	10.6	0.2	0.0	0.0	0.2				
Mean cores	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0				
no./m2 (box)	50	50	0	0	0	0	50	0	0	0	265	5	0	0	5				
no./m2 (core)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0				
<b>No. species (box)</b>	<b>6</b>																		
<b>No. species (core)</b>	<b>0</b>																		
<b>Total Organisms at Station</b>															<b>85</b>				
<b>Station IM3</b>		Depth -5.50m AHD			56 364693		6332101		Sampled August 15th-18th 2020										
Replicates	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod Nassarius	Gastropod Bedeva	Bivalve Corbula	Bivalve Soletellina	Bivalve Paphia	Bivalve Anadara	Bivalve Cyamiomactra	Bivalve Trichomya	Ophuroid	Barnacle	Fish	Crab				
IM3.1	1	0	0	0	0	0	5	0	0	0	0	0	0	0	0				
IM3.2	3	1	0	0	0	0	10	0	0	0	0	0	0	0	0				
IM3.3	2	0	0	0	0	2	6	0	0	0	0	0	0	0	0				
IM3.4	3	0	0	0	0	0	2	0	0	0	0	0	0	0	0				
IM3.5	2	0	0	0	0	3	7	0	0	0	0	0	0	0	0				
IM3.6	0	1	0	0	0	0	7	0	0	0	0	0	0	0	0				
IM3.7	0	1	0	0	0	0	2	0	0	0	0	0	0	0	0				
Mean/station (boxes)	2.2	0.2	0.0	0.0	0.0	1.0	6.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0				
Mean cores	0.0	1.0	0.0	0.0	0.0	0.0	4.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0				
no./m2 (box)	55	5	0	0	0	25	150	0	0	0	0	0	0	0	0				
no./m2 (core)	0	20	0	0	0	0	90	0	0	0	0	0	0	0	0				
<b>No. species (box)</b>	<b>4</b>																		
<b>No. species (core)</b>	<b>2</b>																		
<b>Total Organisms at Station</b>															<b>47</b>				



**Table 4.** Number of species found at each Station from February 2012 to August 2020

Station	C1	C2	C3	C4	C5	C6	C7	R1	R2	R3	R4
Feb 2012	10	5	5	7				8	8	5	5
Sept 2012	3	6	4	4				6	3	4	5
March 2013	4	5	7	7				6	5	6	5
Sept 2013	6	6	3	7				5	6	5	4
March 2014	4	3	5	5				6	4	5	3
Sept 2014	3	4	4	8				6	5	6	6
March 2015	3	3	5	3				5	3	6	5
Sept 2015	5	4	4	3				5	3	4	6
March 2016	6	4	5	5	5			6	5	6	4
Sept 2016	7	3	6	5	4	8		8	4	5	6
March 2017	2	4	5	3	5	5		4	5	4	5
Sept 2017	4	4	4	4	4	5		4	3	6	5
March 2018	4	4	8	4	4	3	5	7	8	5	4
Sept 2018	3	4	4	6	5	5	5	4	4	5	5
March 2019	6	3	4	4	6	5	3	4	5	7	3
Sept 2019	5	6	5	5	4	5	6	4	3	7	4
March 2020	5	6	6	4	7	3	6	6	6	7	4
August 2020	6	5	4	4	3	5	5	4	5	7	4
Station	R5	R6	R7	R8	R9	R10	R11	IM1	IM2	IM3	IM4
Feb 2012								7	4	4	5
Sept 2012								4	4	3	5
March 2013								7	5	5	5
Sept 2013								4	3	4	5
March 2014	4	3						5	9	4	5
Sept 2014	3	3						5	6	3	6
March 2015	3	3						5	4	4	5
Sept 2015	5	4						5	5	4	4
March 2016	4	4	8					6	6	3	4
Sept 2016	6	7	7	5	8			6	4	6	3
March 2017	4	4	4	3	5			3	4	3	4
Sept 2017	4	4	4	5	4			5	5	5	5
March 2018	6	3	4	3	4	4	4	5	7	3	4
Sept 2018	5	4	6	4	5	4	4	4	8	4	4
March 2019	5	4	4	4	4	6	6	5	5	2	4
Sept 2019	4	4	5	4	4	4	3	6	5	7	5
March 2020	4	4	8	3	4	4	4	7	7	4	4
August 2020	7	5	8	4	5	5	4	5	6	4	6

**Table 5** shows the mean number of marine benthic organisms for each station and species sampled in August 2020 (Sieve boxes only). The table includes depths relative to AHD for each station.

**Table 5 Mean number of marine benthic organisms at Control, Reference and Impact stations**

	Depth (m)	Polychaete thin	Polychaete mud	Polychaete thick	Gastropod <i>Nassarius</i>	Gastropod <i>Bedeva</i>	Bivalve <i>Corbula</i>	Bivalve <i>Soletellina</i>	Bivalve <i>Paphia</i>	Bivalve <i>Anadara</i>	Bivalve <i>Cyamiomactra</i>	Bivalve <i>Trichomya</i>	Ophuroid	Bivalve <i>Saccostrea</i>	Fish	Crab
C1	-4.5	1.6	0.8	0.0	0.0	0.0	3.4	13.2	0.0	0.0	0.2	0.0	0.2	0.0	0.0	0.0
C2	-4.5	3.0	4.0	0.0	0.0	0.0	3.4	6.6	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0
C3	-5.5	2.8	0.6	0.0	0.0	0.0	2.4	3.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
C4	-6.0	1.6	0.8	0.0	0.0	0.0	2.8	17.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
C5	-6.0	4.2	7.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.6	0.0	0.0	0.0
C6	-5.5	1.4	0.2	0.0	0.0	0.0	1.2	2.8	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0
C7	-5.5	2.4	3.2	0.0	0.0	0.0	0.2	0.4	0.0	0.0	0.0	0.0	0.8	0.0	0.0	0.0
R1	-4.5	1.8	0.8	0.0	0.0	0.0	1.0	7.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
R2	-4.5	1.6	0.6	0.0	0.0	0.0	5.0	6.8	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0
R3 (IM5)	-5.5	2.2	0.4	0.0	0.0	0.0	0.2	3.0	0.2	0.0	0.0	17.8	0.0	0.2	0.0	0.0
R4 (IM6)	-6.0	1.0	0.8	0.0	0.0	0.0	1.0	2.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
R5 (IM7)	-5.5	2.0	0.2	0.4	0.0	0.0	2.6	4.0	0.2	0.0	0.4	0.0	0.0	0.0	0.0	0.0
R6	-6.0	2.0	0.4	0.2	0.0	0.0	3.2	13.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
R7	-6.0	1.4	1.0	0.0	0.0	0.0	0.2	2.8	0.0	0.0	0.2	7.4	0.6	0.0	0.0	0.2
R8	-5.5	1.6	0.0	0.0	0.0	0.0	0.4	4.4	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
R9	-4.5	2.4	0.0	0.0	0.0	0.0	3.6	8.6	0.2	0.0	0.2	0.0	0.0	0.0	0.0	0.0
R10	-5.5	2.0	1.6	0.0	0.0	0.0	1.0	3.8	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
R11	-6.0	1.4	3.2	0.0	0.0	0.0	0.0	0.6	0.0	0.0	0.0	0.0	1.4	0.0	0.0	0.0
IM1	-4.5	3.0	0.2	0.0	0.0	0.0	2.6	3.2	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0
IM2	-4.5	2.0	2.0	0.0	0.0	0.0	0.0	2.0	0.0	0.0	0.0	10.6	0.2	0.0	0.0	0.2
IM3	-5.5	2.2	0.2	0.0	0.0	0.0	1.0	6.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
IM4	-6.0	1.6	1.0	0.0	0.0	0.0	0.8	2.6	0.2	0.0	0.0	0.0	0.4	0.0	0.0	0.0

August 2020

## Sediment Analysis

The sediment in the mud basin of Lake Macquarie off Summerland Point, in Chain Valley Bay and Bardens Bay was largely composed of fine grey or black mud that sometimes was mildly plastic in nature (able to be molded into a coherent shape) or very fine and fluid, with a small amount of medium to fine black sand in some samples. Large amounts of large and small shell fragments were also present in the sediment at most stations sampled.

A description of the sediment samples collected in August 2020 is shown in **Table 6**.

**Table 6.** Description of Sediment collected from Sampling Stations in August 2020

Station	Description	Volume (mL)
<b>C1</b>	Fine black mud with some shell	250
<b>C2</b>	Fine black mud with some shell and black sand	250
<b>C3</b>	Fine black mud	250
<b>C4</b>	Fine black mud	250
<b>C5</b>	Fine black mud with some shell and black sand	250
<b>C6</b>	Fine black mud with some shell	250
<b>C7</b>	Fine black mud with some shell and black sand	250
<b>R1</b>	Fine black mud with some shell and black sand	250
<b>R2</b>	Fine black mud with some shell and black sand	250
<b>R3 (IM5)</b>	Fine black mud with some shell	250
<b>R4 (IM6)</b>	Fine black mud with some shell and black sand	250
<b>R5 (IM7)</b>	Fine black mud with some shell	250
<b>R6</b>	Fine black mud with some shell	250
<b>R7</b>	Fine black mud with some shell and black sand	250
<b>R8</b>	Fine black mud with some shell and black sand	250
<b>R9</b>	Fine black mud with some shell	250
<b>R10</b>	Fine black mud with some shell	250
<b>R11</b>	Fine black mud with some shell	250
<b>IM1</b>	Fine black mud with some shell	250
<b>IM2</b>	Fine black mud with some shell	250
<b>IM3</b>	Fine black mud with some shell	250
<b>IM4</b>	Fine black mud with some shell	250

**Table 7** shows the percentage of silt in the sediment at each station from February 2012 to August 2020.

**Table 7.** Percent mud in sediment from each station – February 2012 to August 2020.

	<b>Feb 2012</b>	<b>Sept. 2012</b>	<b>Mar. 2013</b>	<b>Sept 2013</b>	<b>Mar. 2014</b>	<b>Sept. 2014</b>	<b>Mar. 2015</b>	<b>Sept. 2015</b>	<b>March 2016</b>
	<b>% Mud</b>	<b>% Mud</b>	<b>% Mud</b>	<b>% Mud</b>	<b>% Mud</b>	<b>% Mud</b>	<b>% Mud</b>	<b>% Mud</b>	<b>% Mud</b>
<b>C1</b>	92	76	60	100	80	72	71	80	96
<b>C2</b>	92	80	60	96	76	77	75	100	96
<b>C3</b>	80	76	64	100	80	75	81	74	96
<b>C4</b>	80	78	64	55	80	75	81	70	96
<b>C5</b>									84
<b>C6</b>									
<b>C7</b>									
<b>R1</b>	80	80	88	40	60	82	80	80	96
<b>R2</b>	94	80	88	60	92	84	97	80	99
<b>R3 IM5</b>	80	82	72	80	88	87	84	80	100
<b>R4 IM6</b>	88	80	80	68	60	85	94	80	100
<b>R5 IM7</b>					80	75	88	65	100
<b>R6</b>					76	78	79	76	100
<b>R7</b>									
<b>R8</b>									
<b>R9</b>									
<b>R10</b>									
<b>R11</b>									
<b>IM1</b>	86	72	68	84	60	77	79	44	100
<b>IM2</b>	80	75	45	60	70	87	78	80	96
<b>IM3</b>	96	75	53	65	72	77	78	75	100
<b>IM4</b>	80	73	56	64	55	86	85	79	98

	Sep-16	Mar-17	Sep-17	Mar-18	Sep-18	Mar-19	Sep-19	Mar-20	Aug-20
	% Mud								
<b>C1</b>	90	68	94	80	80	80	100	94	90
<b>C2</b>	80	92	80	80	64	70	100	86	95
<b>C3</b>	90	80	100	92	80	100	100	96	100
<b>C4</b>	75	98	100	80	40	80	100	92	100
<b>C5</b>	90	92	80	92	64	60	80	29	80
<b>C6</b>	80	92	100	84	60	70	100	100	95
<b>C7</b>				80	60	80	100	29	90
<b>R1</b>	70	96	100	80	80	80	96	96	70
<b>R2</b>	90	80	92	80	84	84	80	80	80
<b>R3 (IM5)</b>	90	96	100	80	96	92	80	92	95
<b>R4 (IM6)</b>	90	96	100	78	92	80	84	98	75
<b>R5 (IM7)</b>	90	88	100	70	80	80	100	94	100
<b>R6</b>	98	98	80	80	78	96	100	92	85
<b>R7</b>	90	94	92	50	80	98	84	92	95
<b>R8</b>	80	98	100	80	82	92	100	92	90
<b>R9</b>				80	84	70	100	80	95
<b>R10</b>				80	96	80	84	80	80
<b>R11</b>				80	30	50	92	100	90
<b>IM1</b>	90	76	96	80	60	80	80	70	95
<b>IM2</b>	98	98	98	92	70	60	80	80	80
<b>IM3</b>	99	96	100	92	96	80	92	90	95
<b>IM4</b>	99	84	92	100	80	80	92	92	90

## Water Quality Profiles - August 2020

At each station, a water quality profile was taken using a calibrated Yeo-Kal 618RU Analyser. Measurements were taken at the surface and at 0.5m intervals to the lakebed. Units of measurement were: Temperature (TEMP) - degrees Celsius; Conductivity (COND) - mS/cm; Salinity (SAL) - parts per thousand; pH; Dissolved Oxygen - % saturation and mg/L; and Turbidity (TURB) - NTU (**Table 8, Appendix 1**).

Up until recently, little significant rain has fallen in the catchments of Lake Macquarie. Annual rainfall in the Cooranbong (Lake Macquarie AWS) region was 839.8 mm in 2017; 859.8 mm in 2018 and 763.4 mm in 2019 (BOM Station Number 061412). The lack of rainfall caused the salinity of the water column to become very high (over 39 parts per thousand by March 2019) and almost uniform from surface to bottom.

The Lake Macquarie region has since received heavy rainfall in August (111.2 mm) and September (64.8 mm) 2019 and again in January (79.6 mm), February (335.4 mm), March (173.0 mm) and July (184.0 mm) 2020 (BOM Station Number 061412). This rainfall lowered the salinity of water in the lake to around 36 parts per thousand in 2019, 33 parts per thousand in March 2020, and 32 parts per thousand in August 2020. Water temperature, salinity, conductivity and pH were found to be uniform from surface to bottom (**Table 8; Appendix 1**).

In August 2020, salinity ranged from 31.80 ppt to 32.36 ppt and conductivity ranged from 596 mS/cm to 702 mS/cm (**Appendix 1**). pH was between 6.70 and 9.93 and water temperature ranged from 14.23°C to 16.15°C (**Appendix 1**). Dissolved oxygen ranged from 80.8% saturation to 111.3 % saturation (**Table 8, Appendix 1**). Throughout the study area, dissolved oxygen concentrations of the bottom water exceeded 100% saturation (**Table 8**).

**Table 8** Physical characteristics of the bottom water - 25<sup>th</sup> August 2020

Station	Temperature °C	Conductivity mS/cm	Salinity ppt	Dissolved Oxygen % sat	Dissolved Oxygen mg/L	pH	Turbidity NTU
C1	15.05	684	32.33	105.1	10.26	8.70	4.1
C2	15.12	610	32.49	110.1	10.49	9.09	5.3
C3	15.03	610	32.53	112.1	10.52	9.27	9.1
C4	14.95	630	32.34	109.8	10.47	9.10	3.9
C5	14.65	611	32.44	117.3	10.52	9.78	5.0
C6	14.85	599	32.30	109.6	10.47	9.10	3.9
C7	15.66	608	32.29	117.9	10.66	9.64	3.1
R1	15.01	702	32.31	102.5	9.87	8.49	2.3
R2	14.60	667	32.11	108.5	10.34	9.07	3.6
R3 (IM5)	15.16	638	32.62	112.4	10.57	9.26	4.6
R4 (IM6)	15.18	640	32.65	111.3	10.50	9.17	4.4
R5 (IM7)	15.03	613	32.37	111.1	10.61	9.19	3.7
R6	14.98	613	32.51	109.9	10.61	9.10	4.1
R7	15.08	607	32.60	118.0	10.55	9.74	2.9
R8	14.89	654	32.35	110.6	10.47	9.18	4.5
R9	14.77	658	32.21	111.0	10.44	9.24	3.0
R10	15.59	608	32.26	119.4	10.72	9.78	3.4

<b>R11</b>	14.91	610	32.57	115.8	10.51	9.59	3.1
<b>IM1</b>	15.03	674	32.34	106.2	10.31	8.79	3.1
<b>IM2</b>	14.97	646	32.42	109.4	10.47	9.06	3.7
<b>IM3</b>	14.92	649	32.36	111.3	10.48	9.23	3.2
<b>IM4</b>	15.01	643	32.48	113.1	10.48	9.36	3.5

## Analysis of Data

### Statistics

Principal component (PC) biplots or multivariate scatterplots produced by the R-statistical program were used to explore the relationship between benthos study sites, animal species found in the sediment, and water quality variables at the lake bed. Points in the matrix were obtained by standardizing the data by subtracting the variable (column) mean from the species (cell) mean and dividing the subsequent value by the variable or column mean (Gabriel, 1971; Gabriel and Odoroff, 1990).

### Biplots

A biplot is a particular kind of scatterplot used for displaying multivariate data which results from mapping a matrix of field observations,  $\mathbf{X}$ , into a 2-dimensional graphical display. The name derives from the fact that this is a *joint* display of the rows and columns of  $\mathbf{X}$ . Sample units (rows) are shown by points and variables (columns) by arrows. Biplots have several appealing properties. Firstly, they are capable of presenting graphically large amounts of information on composition, structure and relationships with surpassing ease and efficiency. It enables a truly global look at the data.

### Interpretation of Biplots

#### Sample Points

- The proximity of any pair of sample points is directly proportional to their resemblance with respect to all the variables studied, the closer the points the greater the resemblance;
- Points close to the origin tend to be representative of the sample as a whole, that is, they tend to be average samples,
- Points far from the origin are atypical in that they possess usually large or small values of one or more variables.

### Variable Arrows


- The origin of the configuration of arrows marks the mean value of each variable, an important reference point.
- Arrows can be extended through the origin (by eye) in either direction to any desired extent.
- With increasing distance from the origin along an arrow in the direction of an arrow, the value of the variable increases steadily above its mean; similarly, with increasing distance from the origin along an arrow extended by eye in the opposite direction, the value of a variable falls increasingly below its mean.
- Arrow length is directly proportional to the correlation coefficient,  $r$ , between the two variables. The smaller the angle the stronger the correlation. Variables  $x$  and  $y$  with arrows subtending an angle of:
  1.  $0^\circ$  are perfectly correlated  $r_{xy} = 1$
  2.  $90^\circ$  are strictly uncorrelated  $r_{xy} = 0$
  3.  $0^\circ \leq \text{Angle} < 90^\circ$   $0 \leq r_{xy} < 1$
  4.  $90^\circ \geq \text{Angle} \leq 180^\circ$   $0 < r_{xy} < -1$

From 3 it follows that variables whose arrows subtend angles less than  $90^\circ$  are positively correlated, and from 4, that variables whose arrows subtend angles greater than  $90^\circ$  are negatively correlated; in particular, where the angle is  $180^\circ$ ,  $r_{xy} = -1$ .

In general, long arrows can be regarded as more useful in interpretation than short arrows. They have greater influence in differentiating sites.

### **Relationship between benthic organisms and stations**

**Figure 5** shows a biplot representing the relationship between marine benthic organisms and stations for the August 2020 survey.



**Figure 5.** Relationship between benthic organisms and sampling stations – Lake Macquarie benthos survey August 2020 (PC biplot goodness-of-fit: 75.14%)

Station		Organism
C1 – Control Station C1	R8 – Reference Station R8	S1 <i>Sthenelais pettiboneae</i>
C2 – Control Station C2	R9 – Reference Station R9	S2 Polychaete mud
C3 – Control Station C3	R10 – Reference Station R10	S3 Polychaete thick
C4 – Control Station C4	R11 – Reference Station R11	S6 <i>Corbula truncata</i>
C5 – Control Station C5	IM1 – Impact Station IM1	S7 <i>Soletellina alba</i>
C6 – Control Station C6	IM2 – Impact Station IM2	S8 <i>Paphia undulata</i>
C7 – Control Station C7	IM3 – Impact Station IM3	S9 <i>Cyamiomactra</i>
R1 – Reference Station R1	IM4 – Impact Station IM4	S11 Ophuroid (Brittle stars)
R2 – Reference Station R2	IM5 – Impact Station IM5	S12 <i>Saccostrea</i> (Oysters)
R6 – Reference Station R6	IM6 – Impact Station IM6	S13 <i>Trichomya hirsuta</i> (mussels)
R7 – Reference Station R7	IM7 – Impact Station IM7	

Five species differentiated sampling stations during the August 2020 sampling period:

- The bivalve *Corbula truncata* (S6) defined the Control Stations C1-C4 and the Reference Stations R2, R6 and R9.
- The bivalve *Soletellina alba* (S7) differentiated the Control Stations C1, C3, C4 and C6; the Reference Stations R1, R2, R6 and R9; and the Impact Stations IM1, IM3 and IM7.
- The Polychaete mud worm designated S2 and brittle stars (S11), characterized stations R11 and C5.
- The mussel species *Trichomya hirsuta* was found at the Impact Stations IM2, IM5 and R7.

## Conclusions

The results from the August 2020 benthic communities monitoring results show compliance to the Schedule 4 Environmental Conditions - underground mining of SSD5465 - Modification 2 in the Performance Measures table with respect to the Subsidence Impact Performance Measure for Benthic communities which display nil to minor environmental consequences due to underground mining.

The below summary of findings outlines the historical basis for this compliance statement and the compliance is detailed in the table below.

Conditions from SSD-5465 – Mod 3	Compliance Status and Comments
Schedule 4 Environmental Conditions – underground mining Performance Measures – Natural Environment Biodiversity – Benthic Communities  Subsidence Impact Performance Measure – Minor environmental consequences, including minor changes composition and/or distribution.	Compliant – See section 16 - Conclusions
Measurements undertaken by generally accepted methods.  Measures Methods fully described.	Compliant – See section 4 and 5  Compliant – See section 4 and 5

In August 2020, 22 benthic stations were sampled in the study area. A total of 1367 organisms greater than 1mm in size were found, comprising 11 species. This compares with the results from March and

September 2019 where 832 and 815 organisms respectively were recorded representing approximately eleven species. As in previous years, polychaete worms and bivalve molluscs were the most frequently encountered animals. Stations were distinguished by the relative abundance of the dominant species. Water depth was not in any way important in determining the species composition at a station.

Physical variables such as salinity, conductivity and turbidity of the bottom water had little influence on the species composition of the benthos. Dissolved oxygen concentration, however, can have a major effect on abundance. It is clear that major extinction events have occurred in the mud basin of Lake Macquarie. The evidence for this lies in the presence of large numbers of intact but dead bivalve shells entombed in the mud. The cause of extinction events appears to be prolonged dissolved oxygen depletion of bottom water. Prolonged dissolved oxygen depletion of the bottom water was measured during the water quality study conducted by Laxton and Laxton (1983 to 1997). Low concentrations of dissolved oxygen in the bottom water were also recorded during the March 2020 sampling period. The dissolved oxygen levels of the bottom waters of Lake Macquarie are currently over 100% saturation.

Bottom sediment in the study area was composed of fine black mud with varying proportions of black sand and shell fragments. In March 2020 some changes to the composition of the upper 100mm of the bottom sediments were detected. At Stations C5 and C7 the sediment comprised mostly sand where previously it was fine black silt. In August 2020, sediment comprised mostly of fine black silt with some shell.

These results appear to support the notion that increasing the water depth by the predicted 0.8m subsidence has, to date, had no discernible effect on the composition and abundance of organisms making up the benthos of the mud basin.

## References

Gabriel, K.R. (1971) The biplot graphical display of matrices with applications to principal component analysis. *Biometrika* 58: 453-525.

Gabriel, K.R. and Odoroff, C.L. (1990) Biplots in biomedical research. *Statistics in Medicine* 9: 469-485.

Laxton, J.H. and Emma Laxton (2007). Aquatic Biology of Chain Valley Bay Lake Macquarie, NSW. Report to Peabody/Lake Coal Chain Valley Colliery.

## Acknowledgements

We wish to acknowledge the help of Mr Chris Ellis, Mr Wade Covey, Mr Chris Armit and Mr Lachlan McWha in facilitating the study.

# Appendix 1

## Water Quality Profile Results for August 2020

Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)
C1	25/08/2020	14:20:24	0.20	14.85	8000	675	2.30	8.42	31.92	101.10	10.13
	25/08/2020	14:20:56	0.56	14.84	8000	678	1.00	8.57	31.93	102.90	10.15
	25/08/2020	14:21:21	1.07	14.80	8000	680	1.00	8.70	31.94	104.40	10.16
	25/08/2020	14:21:46	1.53	14.73	8000	681	1.10	8.76	31.97	105.00	10.17
	25/08/2020	14:22:08	1.99	14.65	8000	682	1.10	8.80	32.02	105.30	10.17
	25/08/2020	14:22:30	2.56	14.56	8000	683	1.20	8.86	32.04	105.90	10.17
	25/08/2020	14:22:47	3.07	14.55	8000	684	1.20	8.91	32.05	106.40	10.16
	25/08/2020	14:23:13	3.50	14.62	8000	684	1.30	8.95	32.10	107.10	10.18
	25/08/2020	14:23:36	4.07	15.02	8000	685	2.10	8.61	32.31	104.00	10.19
	25/08/2020	14:23:53	4.53	15.04	8000	685	3.20	8.57	32.33	103.60	10.21
	25/08/2020	14:25:14	4.57	15.05	8000	684	4.10	8.70	32.33	105.10	10.26
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)
C2	25/08/2020	16:03:58	0.20	15.96	8000	601	2.20	9.08	32.27	111.70	10.62
	25/08/2020	16:04:09	0.51	15.94	8000	603	2.30	9.21	32.25	113.30	10.61
	25/08/2020	16:04:22	1.08	15.57	8000	604	2.40	9.29	32.24	113.50	10.59
	25/08/2020	16:04:35	1.46	15.18	8000	606	2.50	9.26	32.30	112.20	10.56
	25/08/2020	16:04:49	1.88	14.93	8000	606	2.50	9.33	32.28	112.40	10.53
	25/08/2020	16:05:05	1.94	14.91	8000	607	2.60	9.35	32.28	112.70	10.52
	25/08/2020	16:05:24	2.61	14.74	8000	608	2.70	9.47	32.30	113.80	10.50
	25/08/2020	16:05:34	3.19	14.70	8000	608	2.70	9.49	32.28	113.90	10.49
	25/08/2020	16:05:45	3.38	14.68	8000	608	2.80	9.55	32.28	114.50	10.49
	25/08/2020	16:05:53	3.99	14.60	8000	609	2.90	9.61	32.30	115.10	10.48
	25/08/2020	16:06:06	4.36	14.61	8000	609	3.30	9.61	32.31	115.10	10.48
	25/08/2020	16:06:25	4.82	15.12	8000	610	5.30	9.08	32.49	110.10	10.49
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)
C3	25/08/2020	15:56:58	0.25	15.70	8000	600	2.10	8.87	32.26	108.60	10.58
	25/08/2020	15:57:15	0.54	15.64	8000	603	2.40	8.92	32.27	109.10	10.58
	25/08/2020	15:57:28	0.99	15.49	8000	604	2.40	9.01	32.27	109.80	10.58
	25/08/2020	15:57:42	1.48	15.24	8000	606	2.50	9.15	32.27	111.00	10.57
	25/08/2020	15:57:56	2.02	14.79	8000	607	2.60	9.30	32.26	111.80	10.55
	25/08/2020	15:58:10	2.54	14.77	8000	607	2.70	9.37	32.26	112.60	10.53
	25/08/2020	15:58:23	3.00	14.59	8000	608	2.60	9.53	32.32	114.10	10.52
	25/08/2020	15:58:46	3.48	14.58	8000	609	2.80	9.48	32.29	113.50	10.50
	25/08/2020	15:58:57	4.08	14.54	8000	609	2.80	9.54	32.29	114.10	10.50
	25/08/2020	15:59:08	4.51	14.48	8000	610	3.00	9.78	32.30	116.80	10.49
	25/08/2020	15:59:24	5.09	14.50	8000	610	3.10	9.68	32.30	115.70	10.47
	25/08/2020	15:59:40	5.56	15.03	8000	610	9.10	9.27	32.53	112.10	10.52
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)
C4	25/08/2020	15:37:19	0.23	15.60	8000	619	2.10	8.97	32.15	109.50	10.53
	25/08/2020	15:37:31	0.54	15.61	8000	622	2.40	8.98	32.15	109.70	10.53
	25/08/2020	15:38:08	1.00	15.54	8000	625	2.70	9.09	32.15	110.90	10.54
	25/08/2020	15:38:28	1.50	15.56	8000	625	2.70	9.18	32.17	112.00	10.55
	25/08/2020	15:39:02	2.03	15.53	8000	626	2.70	9.31	32.17	113.50	10.55
	25/08/2020	15:39:15	2.44	15.33	8000	627	3.00	9.35	32.28	113.70	10.53
	25/08/2020	15:39:28	2.93	15.07	8000	628	2.80	9.48	32.25	114.60	10.52
	25/08/2020	15:39:42	3.52	14.96	8000	628	2.90	9.36	32.27	112.90	10.49
	25/08/2020	15:39:56	3.97	14.97	8000	629	2.90	9.25	32.28	111.50	10.48
	25/08/2020	15:40:13	4.41	14.94	8000	629	3.60	9.20	32.33	111.00	10.47
	25/08/2020	15:40:27	4.79	14.95	8000	630	4.10	9.09	32.32	109.60	10.46
	25/08/2020	15:40:41	5.09	14.95	8000	630	3.90	9.10	32.34	109.80	10.47

Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)
C5	25/08/2020	16:35:33	0.20	15.24	8000	602	2.20	9.08	32.35	110.10	10.57
	25/08/2020	16:35:47	0.56	15.01	8000	605	2.40	9.19	32.38	111.00	10.57
	25/08/2020	16:35:59	1.05	14.98	8000	606	2.50	9.19	32.37	110.90	10.56
	25/08/2020	16:36:16	1.51	14.96	8000	607	2.50	9.34	32.37	112.70	10.56
	25/08/2020	16:36:28	2.06	14.96	8000	607	3.20	9.26	32.38	111.70	10.56
	25/08/2020	16:36:42	2.49	14.96	8000	608	2.60	9.32	32.38	112.50	10.56
	25/08/2020	16:36:58	3.02	14.95	8000	608	2.70	9.41	32.38	113.50	10.56
	25/08/2020	16:37:16	3.55	14.93	8000	609	2.80	9.50	32.37	114.60	10.56
	25/08/2020	16:37:30	4.07	14.97	8000	609	2.70	9.57	32.40	115.50	10.56
	25/08/2020	16:37:43	4.52	14.79	8000	609	2.80	9.68	32.37	116.40	10.55
	25/08/2020	16:37:59	5.06	14.75	8000	610	2.80	9.78	32.40	117.50	10.54
	25/08/2020	16:38:15	5.53	14.68	8000	610	2.80	9.72	32.40	116.70	10.54
	25/08/2020	16:38:27	6.06	14.64	8000	610	2.90	9.83	32.41	117.90	10.53
	25/08/2020	16:38:43	6.54	14.71	8000	611	3.20	9.80	32.44	117.80	10.53
	25/08/2020	16:38:54	7.10	14.65	8000	611	5.00	9.78	32.44	117.30	10.52
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)
C6	25/08/2020	15:45:24	0.23	15.60	8000	616	3.00	8.98	32.14	109.60	10.53
	25/08/2020	15:45:38	0.56	15.61	8000	619	2.40	9.04	32.14	110.40	10.54
	25/08/2020	15:45:51	1.02	15.61	8000	620	2.60	9.04	32.15	110.40	10.55
	25/08/2020	15:46:06	1.50	15.60	8000	621	2.60	9.20	32.15	112.30	10.55
	25/08/2020	15:46:19	2.06	15.57	8000	621	2.70	9.18	32.16	112.00	10.55
	25/08/2020	15:46:35	2.49	15.50	8000	622	2.70	9.25	32.17	112.70	10.55
	25/08/2020	15:46:57	3.02	15.05	8000	623	2.80	9.41	32.30	113.70	10.53
	25/08/2020	15:47:08	3.53	14.90	8000	624	2.90	9.44	32.28	113.80	10.50
	25/08/2020	15:47:29	4.00	14.85	8000	625	3.30	9.20	32.30	110.70	10.49
	25/08/2020	15:48:20	4.62	14.85	8000	599	3.90	9.10	32.30	109.60	10.47
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)
C7	25/08/2020	17:02:04	0.23	16.07	8000	597	2.00	9.09	32.24	112.20	10.72
	25/08/2020	17:02:17	0.55	16.08	8000	600	2.20	9.15	32.24	112.90	10.72
	25/08/2020	17:02:43	1.12	16.07	8000	603	2.50	9.24	32.25	113.90	10.72
	25/08/2020	17:02:58	1.52	16.06	8000	604	2.50	9.27	32.23	114.30	10.72
	25/08/2020	17:03:12	2.04	16.03	8000	605	2.50	9.27	32.23	114.20	10.71
	25/08/2020	17:03:22	2.53	16.01	8000	605	2.60	9.33	32.23	114.90	10.70
	25/08/2020	17:03:34	3.19	15.99	8000	606	2.60	9.36	32.23	115.20	10.70
	25/08/2020	17:03:46	3.37	15.99	8000	606	2.60	9.38	32.23	115.50	10.69
	25/08/2020	17:03:56	3.99	15.98	8000	606	2.60	9.54	32.23	117.40	10.69
	25/08/2020	17:04:10	4.44	15.89	8000	607	2.60	9.56	32.26	117.50	10.68
	25/08/2020	17:04:25	4.76	15.66	8000	608	3.10	9.64	32.29	117.90	10.66
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)
R1	25/08/2020	14:08:22	0.33	15.09	8000	692	0.20	6.70	31.88	80.80	9.80
	25/08/2020	14:09:00	1.01	15.11	8000	695	0.30	7.64	31.89	92.20	9.78
	25/08/2020	14:09:35	1.53	14.99	8000	697	0.20	8.10	31.88	97.50	9.78
	25/08/2020	14:10:11	2.00	14.23	8000	699	0.40	8.50	31.85	100.70	9.75
	25/08/2020	14:10:39	2.48	14.50	8000	700	0.50	8.52	31.96	101.70	9.76
	25/08/2020	14:11:05	3.00	14.83	8000	700	0.60	8.69	32.15	104.50	9.82
	25/08/2020	14:12:03	3.66	14.83	8000	701	0.80	8.98	32.19	107.90	9.87
	25/08/2020	14:12:36	4.12	15.01	8000	702	2.30	8.49	32.31	102.50	9.87

Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)
R2	25/08/2020	14:38:54	0.19	14.81	8000	655	1.20	8.81	31.80	105.60	10.37
	25/08/2020	14:39:13	0.51	14.82	8000	659	1.30	8.97	31.80	107.50	10.37
	25/08/2020	14:39:36	1.01	14.77	8000	661	1.50	9.08	31.81	108.70	10.37
	25/08/2020	14:39:56	1.54	14.68	8000	662	1.50	9.17	31.84	109.60	10.37
	25/08/2020	14:40:11	1.51	14.57	8000	663	1.60	9.20	31.88	109.90	10.37
	25/08/2020	14:40:28	2.01	14.39	8000	664	1.70	9.21	31.94	109.60	10.36
	25/08/2020	14:40:47	2.52	14.33	8000	665	1.70	9.25	31.97	109.90	10.35
	25/08/2020	14:41:06	3.01	14.34	8000	666	1.80	9.26	31.99	110.10	10.35
	25/08/2020	14:41:23	3.49	14.38	8000	666	1.80	9.25	32.01	110.10	10.35
	25/08/2020	14:41:37	4.06	14.55	8000	667	2.30	9.23	32.11	110.30	10.34
	25/08/2020	14:41:53	4.54	14.60	8000	667	3.60	9.07	32.11	108.50	10.34
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)
R3	25/08/2020	15:29:49	0.16	15.63	8000	627	4.10	9.03	32.13	110.40	10.53
	25/08/2020	15:30:02	0.57	15.65	8000	630	2.70	9.01	32.14	110.10	10.52
	25/08/2020	15:30:16	1.05	15.66	8000	631	2.80	9.03	32.14	110.40	10.53
	25/08/2020	15:30:26	1.54	15.65	8000	632	2.80	9.08	32.14	111.00	10.54
	25/08/2020	15:30:41	2.00	15.45	8000	633	2.90	9.21	32.20	112.10	10.54
	25/08/2020	15:30:57	2.51	15.21	8000	634	2.90	9.40	32.18	113.80	10.54
	25/08/2020	15:31:08	2.99	14.92	8000	635	2.90	9.55	32.19	115.00	10.52
	25/08/2020	15:31:21	3.49	14.84	8000	635	3.00	9.58	32.19	115.20	10.50
	25/08/2020	15:31:32	4.02	14.9	8000	636	3.10	9.60	32.25	115.60	10.50
	25/08/2020	15:31:42	4.50	14.93	8000	636	3.00	9.68	32.26	116.60	10.50
	25/08/2020	15:31:53	5.01	14.96	8000	637	3.10	9.65	32.43	116.50	10.49
	25/08/2020	15:32:05	5.48	15.07	8000	637	3.30	9.42	32.55	114.00	10.50
	25/08/2020	15:32:20	6.00	15.12	8000	637	3.50	9.29	32.59	112.70	10.50
	25/08/2020	15:32:43	6.44	15.14	8000	638	3.90	9.26	32.61	112.30	10.51
	25/08/2020	15:33:02	6.86	15.16	8000	638	4.60	9.26	32.62	112.40	10.51
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)
R4	25/08/2020	15:23:05	0.30	15.59	8000	626	2.20	8.79	32.15	107.30	10.53
	25/08/2020	15:23:22	0.55	15.58	8000	632	2.60	8.94	32.15	109.10	10.53
	25/08/2020	15:23:33	1.07	15.52	8000	633	2.60	9.00	32.16	109.70	10.53
	25/08/2020	15:23:48	1.52	15.51	8000	634	2.70	9.08	32.15	110.60	10.53
	25/08/2020	15:24:05	2.05	15.36	8000	635	2.80	9.20	32.17	111.80	10.53
	25/08/2020	15:24:20	2.54	14.90	8000	636	2.80	9.40	32.20	113.20	10.52
	25/08/2020	15:24:38	3.03	14.87	8000	637	2.90	9.47	32.20	114.00	10.50
	25/08/2020	15:24:54	3.56	14.88	8000	638	2.90	9.51	32.20	114.50	10.50
	25/08/2020	15:25:09	4.04	14.83	8000	638	2.90	9.55	32.22	114.80	10.49
	25/08/2020	15:25:27	4.49	14.86	8000	639	2.90	9.55	32.28	115.00	10.49
	25/08/2020	15:25:39	4.96	14.85	8000	639	3.00	9.64	32.32	116.00	10.49
	25/08/2020	15:25:55	5.10	14.87	8000	639	3.10	9.53	32.39	114.90	10.49
	25/08/2020	15:26:05	5.50	15.04	8000	640	3.60	9.43	32.49	114.10	10.49
	25/08/2020	15:26:15	6.02	15.15	8000	640	4.00	9.24	32.63	112.20	10.49
	25/08/2020	15:26:33	6.49	15.18	8000	640	4.40	9.17	32.65	111.30	10.50
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)
R5	25/08/2020	17:32:28	0.18	15.77	8000	603	2.10	9.00	32.15	110.20	10.69
	25/08/2020	17:32:42	0.52	15.77	8000	607	2.30	9.04	32.16	110.70	10.69
	25/08/2020	17:32:53	1.01	15.78	8000	608	2.40	9.08	32.16	111.20	10.69
	25/08/2020	17:33:04	1.51	15.79	8000	609	2.60	9.15	32.16	112.10	10.69
	25/08/2020	17:33:18	2.04	15.79	8000	610	2.60	9.22	32.15	113.00	10.70
	25/08/2020	17:33:29	2.45	15.79	8000	610	2.60	9.24	32.16	113.30	10.70
	25/08/2020	17:33:40	3.12	15.79	8000	610	2.60	9.34	32.16	114.50	10.70
	25/08/2020	17:33:53	3.51	15.80	8000	611	2.70	9.42	32.17	115.50	10.69
	25/08/2020	17:34:03	4.08	15.80	8000	611	2.70	9.44	32.17	115.70	10.69
	25/08/2020	17:34:14	4.53	15.42	8000	612	3.00	9.70	32.27	118.10	10.67
	25/08/2020	17:34:25	5.07	15.07	8000	613	3.60	9.52	32.31	115.10	10.64
	25/08/2020	17:34:40	5.46	15.03	8000	613	3.70	9.19	32.37	111.10	10.61

Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)	
R6	25/08/2020	17:25:21	0.19	15.69	8000	599	2.00	9.05	32.17	110.70	10.72	
	25/08/2020	17:25:35	0.53	15.71	8000	603	2.20	9.11	32.17	111.40	10.71	
	25/08/2020	17:25:47	1.02	15.70	8000	605	2.30	9.20	32.17	112.60	10.71	
	25/08/2020	17:26:04	1.56	15.70	8000	606	2.40	9.15	32.17	111.90	10.71	
	25/08/2020	17:26:17	2.01	15.71	8000	607	2.50	9.22	32.18	112.90	10.71	
	25/08/2020	17:26:29	2.52	15.69	8000	608	2.60	9.26	32.18	113.30	10.71	
	25/08/2020	17:26:42	3.00	15.65	8000	609	2.70	9.48	32.19	115.80	10.71	
	25/08/2020	17:26:56	3.49	15.64	8000	609	2.60	9.44	32.21	115.30	10.71	
	25/08/2020	17:27:08	4.03	15.49	8000	609	2.60	9.59	32.24	116.80	10.70	
	25/08/2020	17:27:23	4.50	15.21	8000	610	2.70	9.73	32.23	117.90	10.69	
	25/08/2020	17:27:33	4.99	15.02	8000	611	3.00	9.93	32.23	119.80	10.67	
	25/08/2020	17:27:47	5.50	14.92	8000	612	3.40	9.75	32.27	117.40	10.62	
	25/08/2020	17:27:59	6.01	15.10	8000	613	4.00	9.26	32.55	112.20	10.60	
	25/08/2020	17:28:25	5.56	14.98	8000	613	4.10	9.10	32.51	109.90	10.61	
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)	
R7	25/08/2020	16:12:36	0.18	15.48	8000	596	2.10	9.02	32.30	109.90	10.57	
	25/08/2020	16:12:49	0.52	15.49	8000	599	2.40	9.01	32.31	109.90	10.56	
	25/08/2020	16:12:59	1.05	15.45	8000	601	2.40	9.10	32.32	110.90	10.56	
	25/08/2020	16:13:13	1.49	15.46	8000	602	2.50	9.10	32.32	110.90	10.55	
	25/08/2020	16:13:33	2.10	15.32	8000	603	2.60	9.31	32.36	113.10	10.56	
	25/08/2020	16:13:57	3.05	14.90	8000	605	2.70	9.50	32.30	114.50	10.55	
	25/08/2020	16:14:10	3.51	14.81	8000	605	2.70	9.62	32.30	115.70	10.54	
	25/08/2020	16:14:24	4.00	14.81	8000	606	2.80	9.65	32.30	116.00	10.53	
	25/08/2020	16:14:35	4.42	14.94	8000	606	2.80	9.62	32.34	116.00	10.52	
	25/08/2020	16:14:43	5.01	15.06	8000	606	2.90	9.54	32.59	115.60	10.53	
	25/08/2020	16:14:54	5.35	15.06	8000	607	2.90	9.62	32.58	116.40	10.55	
	25/08/2020	16:15:06	5.99	15.06	8000	607	2.90	9.62	32.60	116.50	10.55	
	25/08/2020	16:15:18	6.44	15.08	8000	607	2.90	9.74	32.60	118.00	10.55	
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)	
R8	25/08/2020	14:53:54	0.26	15.07	8000	643	1.90	8.65	32.10	104.50	10.49	
	25/08/2020	14:54:15	0.55	15.07	8000	647	2.20	8.84	32.11	106.80	10.49	
	25/08/2020	14:54:29	1.10	15.06	8000	648	2.30	8.88	32.12	107.20	10.49	
	25/08/2020	14:54:47	1.49	15.03	8000	649	2.40	8.95	32.15	108.00	10.50	
	25/08/2020	14:55:04	1.98	15.02	8000	650	2.40	9.01	32.16	108.70	10.50	
	25/08/2020	14:55:28	2.49	15.00	8000	651	2.50	9.10	32.18	109.70	10.50	
	25/08/2020	14:55:47	2.89	15.03	8000	651	2.40	9.11	32.26	110.00	10.50	
	25/08/2020	14:56:07	3.44	14.75	8000	652	2.60	9.24	32.24	110.90	10.48	
	25/08/2020	14:56:25	3.95	14.69	8000	653	2.70	9.26	32.24	111.00	10.47	
	25/08/2020	14:56:49	4.54	14.73	8000	653	2.80	9.19	32.26	110.40	10.47	
	25/08/2020	14:57:03	5.08	14.90	8000	653	2.80	9.18	32.34	110.70	10.48	
	25/08/2020	14:57:20	5.52	14.89	8000	654	3.70	9.22	32.35	111.10	10.47	
	25/08/2020	14:57:40	5.74	14.89	8000	654	4.50	9.18	32.35	110.60	10.47	
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)	
R9	25/08/2020	14:46:28	0.15	14.84	8000	648	1.80	8.96	31.81	107.50	10.45	
	25/08/2020	14:46:48	0.55	14.86	8000	652	2.00	9.10	31.81	109.30	10.45	
	25/08/2020	14:47:00	1.03	14.87	8000	653	2.20	9.10	31.81	109.20	10.45	
	25/08/2020	14:47:19	1.51	14.90	8000	654	2.20	9.24	31.84	111.00	10.46	
	25/08/2020	14:47:38	2.09	14.97	8000	655	2.30	9.27	31.90	111.60	10.46	
	25/08/2020	14:47:48	2.52	14.81	8000	656	2.30	9.28	32.00	111.40	10.46	
	25/08/2020	14:48:05	2.99	14.75	8000	657	2.40	9.29	32.11	111.40	10.45	
	25/08/2020	14:48:21	3.50	14.73	8000	658	2.70	9.25	32.20	111.00	10.45	
	25/08/2020	14:48:36	3.99	14.77	8000	658	3.00	9.24	32.21	111.00	10.44	

Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)	
R10	25/08/2020	17:14:20	0.23	16.06	8000	597	2	8.94	32.17	110.2	10.71	
	25/08/2020	17:14:31	0.52	16.08	8000	600	2.1	8.96	32.18	110.4	10.71	
	25/08/2020	17:14:45	1.01	16.11	8000	602	2.2	9	32.18	111	10.71	
	25/08/2020	17:14:57	1.52	16.13	8000	604	2.3	9.05	32.18	111.7	10.72	
	25/08/2020	17:15:09	2	16.12	8000	605	2.4	9.2	32.18	113.5	10.72	
	25/08/2020	17:15:22	2.5	16.15	8000	605	2.5	9.19	32.19	113.5	10.72	
	25/08/2020	17:15:35	3.06	16.05	8000	606	2.5	9.32	32.18	114.9	10.72	
	25/08/2020	17:15:46	3.49	16.03	8000	606	2.5	9.42	32.19	116	10.72	
	25/08/2020	17:15:57	3.99	16.04	8000	607	2.6	9.46	32.2	116.6	10.72	
	25/08/2020	17:16:09	4.54	15.77	8000	608	2.6	9.6	32.25	117.7	10.72	
	25/08/2020	17:16:23	4.62	15.59	8000	608	3.4	9.78	32.26	119.4	10.72	
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)	
R11	25/08/2020	16:23:41	0.29	15.29	8000	598	2	9.07	32.33	110.1	10.57	
	25/08/2020	16:23:54	0.52	15.12	8000	602	2.3	9.15	32.36	110.8	10.54	
	25/08/2020	16:24:07	1.03	14.79	8000	604	2.4	9.27	32.32	111.4	10.54	
	25/08/2020	16:24:25	1.57	14.72	8000	605	2.5	9.27	32.37	111.3	10.52	
	25/08/2020	16:24:38	2.07	14.78	8000	606	2.5	9.26	32.43	111.4	10.52	
	25/08/2020	16:24:48	2.66	14.97	8000	606	2.5	9.29	32.5	112.2	10.53	
	25/08/2020	16:25:01	3.04	14.98	8000	607	2.6	9.39	32.5	113.5	10.54	
	25/08/2020	16:25:15	3.52	14.75	8000	607	2.7	9.56	32.48	114.9	10.52	
	25/08/2020	16:25:30	4	14.76	8000	608	2.7	9.48	32.48	114	10.51	
	25/08/2020	16:25:44	4.39	14.79	8000	608	2.9	9.48	32.5	114.1	10.51	
	25/08/2020	16:25:57	4.95	14.84	8000	609	2.9	9.37	32.53	112.9	10.5	
	25/08/2020	16:26:12	5.43	14.85	8000	609	3	9.37	32.54	113	10.5	
	25/08/2020	16:26:33	5.94	14.88	8000	609	3.2	9.42	32.56	113.6	10.5	
	25/08/2020	16:27:07	6.08	14.91	8000	610	3.1	9.59	32.57	115.8	10.51	
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)	
IM1	25/08/2020	14:29:23	0.19	14.76	8000	665	0.90	8.53	31.91	102.20	10.30	
	25/08/2020	14:29:45	0.52	14.74	8000	668	1.10	8.74	31.92	104.70	10.29	
	25/08/2020	14:30:03	1.01	14.74	8000	669	1.20	8.85	31.92	106.00	10.30	
	25/08/2020	14:30:25	1.50	14.67	8000	670	1.30	8.97	31.96	107.40	10.30	
	25/08/2020	14:30:44	1.99	14.64	8000	671	1.40	9.02	32.02	107.90	10.30	
	25/08/2020	14:31:01	2.49	14.70	8000	672	1.50	9.04	32.10	108.30	10.30	
	25/08/2020	14:31:19	2.95	14.72	8000	673	1.50	9.07	32.12	108.70	10.30	
	25/08/2020	14:31:41	3.52	14.75	8000	673	1.60	9.13	32.15	109.60	10.30	
	25/08/2020	14:32:08	4.01	15.00	8000	674	1.90	8.80	32.32	106.30	10.30	
	25/08/2020	14:32:31	4.50	15.03	8000	674	3.10	8.79	32.34	106.20	10.31	
Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)	
IM2	25/08/2020	15:07:38	0.18	15.21	8000	638	2.40	8.95	32.15	108.40	10.50	
	25/08/2020	15:07:53	0.51	15.21	8000	640	2.40	9.05	32.15	109.60	10.50	
	25/08/2020	15:08:10	1.01	15.19	8000	641	2.50	9.08	32.15	109.90	10.51	
	25/08/2020	15:08:25	1.50	15.18	8000	641	2.60	9.17	32.15	111.00	10.51	
	25/08/2020	15:08:46	2.05	15.10	8000	642	2.60	9.21	32.17	111.30	10.50	
	25/08/2020	15:09:07	2.54	14.89	8000	643	2.70	9.27	32.20	111.60	10.49	
	25/08/2020	15:09:26	3.01	14.86	8000	643	2.70	9.33	32.20	112.30	10.49	
	25/08/2020	15:09:45	3.51	14.72	8000	644	2.80	9.41	32.22	112.80	10.48	
	25/08/2020	15:10:11	3.98	14.68	8000	644	2.80	9.43	32.22	113.10	10.47	
	25/08/2020	15:10:51	4.45	14.88	8000	645	3.10	9.12	32.32	109.80	10.46	
	25/08/2020	15:11:49	5.21	14.97	8000	646	3.70	9.06	32.42	109.40	10.47	

Station	Date	Time	Depth (M)	Temp (C)	Cond (us/cm)	Cond (ms/cm)	Turb (ntu)	pH (pH)	Sal (ppt)	D.O. (%sat)	D.O. (mg/l)
<b>IM3</b>	25/08/2020	15:01:09	0.23	15.13	8000	640	2.00	8.94	32.14	108.10	10.51
	25/08/2020	15:01:23	0.56	15.11	8000	642	2.20	8.99	32.14	108.70	10.51
	25/08/2020	15:01:38	1.02	15.12	8000	643	2.30	9.01	32.14	109.00	10.52
	25/08/2020	15:01:56	1.54	15.11	8000	644	2.40	9.09	32.15	109.80	10.51
	25/08/2020	15:02:14	1.98	15.01	8000	645	2.40	9.17	32.17	110.60	10.51
	25/08/2020	15:02:38	2.53	14.94	8000	646	2.50	9.24	32.20	111.30	10.50
	25/08/2020	15:02:57	3.04	14.80	8000	647	2.60	9.29	32.22	111.70	10.49
	25/08/2020	15:03:18	3.50	14.72	8000	647	2.60	9.36	32.22	112.30	10.48
	25/08/2020	15:03:35	3.96	14.69	8000	648	2.70	9.40	32.23	112.70	10.48
	25/08/2020	15:03:47	4.42	14.69	8000	648	2.80	9.37	32.24	112.40	10.48
	25/08/2020	15:04:00	4.96	14.89	8000	648	2.80	9.30	32.34	112.10	10.47
	25/08/2020	15:04:15	5.52	14.92	8000	649	3.70	9.26	32.38	111.60	10.47
	25/08/2020	15:04:53	5.23	14.92	8000	649	3.20	9.23	32.36	111.30	10.48
<b>Station</b>	<b>Date</b>	<b>Time</b>	<b>Depth (M)</b>	<b>Temp (C)</b>	<b>Cond (us/cm)</b>	<b>Cond (ms/cm)</b>	<b>Turb (ntu)</b>	<b>pH (pH)</b>	<b>Sal (ppt)</b>	<b>D.O. (%sat)</b>	<b>D.O. (mg/l)</b>
<b>IM4</b>	26/08/2020	15:14:40	0.16	15.47	8000	633	2.30	8.88	32.14	108.10	10.51
	25/08/2020	15:15:44	0.53	15.49	8000	637	2.50	8.99	32.14	109.50	10.51
	25/08/2020	15:16:02	1.01	15.48	8000	638	2.60	9.02	32.15	109.90	10.52
	25/08/2020	15:16:24	1.52	15.47	8000	638	2.70	9.10	32.14	110.80	10.53
	25/08/2020	15:16:41	2.03	15.38	8000	639	2.70	9.19	32.16	111.70	10.52
	25/08/2020	15:16:56	2.50	15.21	8000	640	2.70	9.30	32.16	112.60	10.52
	25/08/2020	15:17:13	3.03	14.92	8000	640	2.80	9.40	32.23	113.30	10.51
	25/08/2020	15:17:26	3.46	14.84	8000	641	2.80	9.48	32.21	114.00	10.50
	25/08/2020	15:17:39	3.99	14.80	8000	641	2.80	9.51	32.21	114.30	10.49
	25/08/2020	15:17:51	4.47	14.88	8000	641	2.80	9.57	32.27	115.20	10.48
	25/08/2020	15:18:06	4.90	14.83	8000	642	2.80	9.78	32.36	117.70	10.49
	25/08/2020	15:18:31	4.94	14.88	8000	642	2.90	9.58	32.31	115.40	10.48
	25/08/2020	15:18:54	5.53	14.93	8000	643	3.10	9.52	32.44	114.90	10.48
	25/08/2020	15:19:20	6.03	15.01	8000	643	3.50	9.36	32.48	113.10	10.48